

# Dragonflies in Devon

**Dave Smallshire**

**January 2026**



Devon is one of the most species-rich counties in the UK for damselflies and dragonflies (Odonata), resulting from its southerly position and wide range of wetland habitats. The maritime effect gives relatively mild winters that limit the amount of freezing in wetlands, although this also cools the springs and summers unless there is a warm airflow from south-east England or the Continent. Habitats range from relatively nutrient-rich lakes, ponds, slow-flowing rivers, canals and ditches of the lowlands to acidic blanket and valley moorland bogs, flushes, streams and fast-flowing rivers of Dartmoor and Exmoor. Most of the important dragonfly sites are at least in part artificial, many resulting from the historic mining of tin, ball clay and China clay. More natural sites include the heathland bogs and springline mires of the moors, the East Devon Pebblebed Heaths and the Blackdown Hills (at which Keeled Skimmer, shown above, is very characteristic). These sites of known importance to dragonflies are listed in the Appendix, with details of their location and the locally- and nationally-important species that they support (or may do). The rest of this document comprises a summary of dragonfly recording in the county followed by a detailed systematic list.

Page 1

## Contents (click on link to go to relevant section)

### History of dragonfly recording in Devon

#### Species accounts

Emerald Damselfly *Lestes sponsa*

Banded Demoiselle *Calopteryx splendens*

Beautiful Demoiselle *Calopteryx virgo*

White-legged Damselfly *Platycnemis pennipes*

Small Red Damselfly *Ceriagrion tenellum*

Southern Damselfly *Coenagrion mercuriale*

Azure Damselfly *Coenagrion puella*

Common Blue Damselfly *Enallagma cyathigerum*

Red-eyed Damselfly *Erythromma najas*

Small Red-eyed Damselfly *Erythromma viridulum*

Blue-tailed Damselfly *Ischnura elegans*

Scarce Blue-tailed Damselfly *Ischnura pumilio*

Large Red Damselfly *Pyrrhosoma nymphula*

Southern Migrant Hawker *Aeshna affinis*

Southern Hawker *Aeshna cyanea*

Brown Hawker *Aeshna grandis*

Common Hawker *Aeshna juncea*

Migrant Hawker *Aeshna mixta*

Norfolk Hawker *Aeshna isosceles*

Vagrant Emperor *Anax ephippiger*

Emperor Dragonfly *Anax imperator*

Lesser Emperor *Anax parthenope*

Hairy Dragonfly *Brachytron pratense*

Common Clubtail *Gomphus vulgatissimus*

Golden-ringed Dragonfly *Cordulegaster boltonii*

Orange-spotted Emerald *Oxygastra curtisii*

Downy Emerald *Cordulia aenea*

Scarlet Darter *Crocothemis erythraea*

Broad-bodied Chaser *Libellula depressa*

Scarce Chaser *Libellula fulva*

Four-spotted Chaser *Libellula quadrimaculata*

Black-tailed Skimmer *Orthetrum cancellatum*

Keeled Skimmer *Orthetrum coerulescens*

Wandering Glider *Pantala flavescens*

Black Darter *Sympetrum danae*

Yellow-winged Darter *Sympetrum flaveolum*

Red-veined Darter *Sympetrum fonscolombii*

Ruddy Darter *Sympetrum sanguineum*

Common Darter *Sympetrum striolatum*

Vagrant Darter *Sympetrum vulgatum*

### APPENDIX: Devon sites for important dragonfly species

## History of dragonfly recording in Devon

Nineteenth century records of dragonflies and damselflies (Odonata) in Devon come from Parfitt (1879), who listed 22 species, although the records of three of these species have not been regarded as authentic in modern times. Since then, records have been collated by the Biological Records Centre's Odonata Recording Scheme (ORS, now the British Dragonfly Society's (BDS) Dragonfly Recording Scheme). For the Devonshire Association's Entomology Group, O. G Watkins collated records in the DA Transactions from 1948 to 1981, but not all of these were considered acceptable for inclusion by the ORS. In particular, a Southern Darter *Sympetrum meridionale* from Dawlish in 1901 was subsequently re-identified by the Natural History Museum as a Common Darter *Sympetrum striolatum*. A "Black-legged Sympetrum (*Sympetrum nigrifemur*)" was recorded at Stover in 1978, but this (sub?)species occurs only in Macaronesia and it is safe to conclude that the record refers to a melanic form of Common Darter.

Common Darters account for seven of the 12 records in the ORS database prior to the First World War, but Red-veined Darter *Sympetrum fonscolombii*, a scarce migrant species, surprisingly featured in 1903 and 1911. The most unusual record for the beginning of the 20<sup>th</sup> century, however, comprised a Vagrant Emperor *Anax ephippiger* in Plymouth on 24 February 1903; a late winter timing has proved to be not unusual for this species, when small incursions, mainly on the south coast, have coincided with depositions of Saharan dust.

Perhaps the most outstanding of all of Devon's dragonfly records comprises three Orange-spotted Emeralds *Oxygastra curtisii* (two males) taken by the River Tamar near Gunnislake by O. G. Watkins in July 1946. This West European endemic was known elsewhere in Britain only from the Moors River, Hampshire, where it was last seen in 1963. There have been many searches along the lower Tamar Valley subsequently, but all have been in vain. There are still eminently suitable calm stretches of the river above a series of weirs, but perhaps the species suffered from toxins released from one of the many mines in the valley. There was also a report of Orange-spotted Emerald from the unlikely location of Braunton Burrows in about 1830, but this was never authenticated.

Bob Merritt, a co-founder of the BDS, lived in Devon for most of 1965-1982. The county survey he initiated in 1977 produced records over the next six years from nearly all of Devon's hectads (10 km × 10 km grid squares). Merritt (1983) included a brief summary of the county's important dragonfly habitats and 5-km square grid maps of the distribution of the 26 species known at the time to be breeding in the county (a substantial increase from the 19 creditable species documented by Parfitt). Merritt corresponded with Watkins, but was reluctant to accept the records of some species following extensive field surveys made during 1965-82.

In 1988, the ORS initiated a project to gather data for important ('key') dragonfly sites. This was pursued in Devon with a concerted 'Key Sites Survey' in 1995, resulting an Inventory of Key Sites for use by countryside agencies, local authorities and land owners and managers (Smallshire, 1996). Sites with recent records of the scarcer species were targeted for surveys, during which breeding evidence was obtained, numbers estimated and the habitat

described, generally during two or three visits per site annually. Site evaluation methods were established, which now form a national standard. Surveys of Key Sites were repeated in 2005/6, producing an increase in the number of sites of national importance (i.e. with healthy breeding populations of Red List species) from eight to 13 and an increase from six to nine confirmed sites of local importance. However, these figures likely represented an increase in recording effort and awareness as much as improvements in the fortunes of dragonflies. Further concerted recording effort around the county was made for a second national atlas in the years up to 2012, the results being published in Cham *et al.*, 2014. More recently, records for the period since 1987 have been analysed for what are now known as Priority Site status: details of these can be found in site inventory files on the BDS website.

Over half (28) of the first 46 records in the ORS database relate to the Bovey Basin and its important dragonfly wetland sites. While Downy Emerald *Cordulia aenea* still has a stronghold here, sadly the Small Red Damselfly *Ceriagrion tenellum* was last seen in 1984, prior to the site's destruction by ball clay extraction. The Bovey Basin remains important for dragonflies today, but the outstanding national importance of Dartmoor and the East Devon Pebblebed Heaths (EDPH) is now more fully appreciated.

The first record of another West European endemic, Southern Damselfly *Coenagrion mercuriale*, came in 1956 from the EDPH, followed by reports from two sites in the Blackdown Hills: Southey Moor in 1959 and Hense Moor in 1962. This is the only species of dragonfly in Devon that has legal protection, initially under the Wildlife and Countryside Act and subsequently following the EU Habitat Regulations. Concerted searches on the northern fringes of Dartmoor during 1995-2005 resulted in the discovery of Southern Damselfly at three further sites, two of which were on the very edges of open moorland, the third being in enclosed pasture. One of the EDPH colonies of this species, at Venn Ottery Common, went extinct around 1990, but was successfully re-established from New Forest stock in 2009. A similar attempt was made at Hense Moor in 2015, but this appears to have failed. The numbers at a long-established colony at Colaton Raleigh Common have declined since 2010 to zero since 2021, for no clear reason. All of the extant sites have been well-monitored by Dave Boyce (for Dartmoor National Park Authority) and RSPB/Lesley Kerry (EDPH sites).

Ruddy Darter *Sympetrum sanguineum* remains strangely scarce in Devon, despite being common in the adjacent counties of Dorset and Somerset. Variable Damselfly *Coenagrion pulchellum* occurs in the grazing marsh ditches of the Somerset Levels but its presence in Devon has never been verified, despite records in the Transactions of the DA from 1879 and 1976-80; this species, especially females, can be easily confused with the widespread Azure Damselfly *C. puella*. The Brown Hawker *Aeshna grandis* is surprisingly rare in Devon, although strangely Parfitt (1879) reported it as "abundant ... in counties including Devonshire, according to Mr Stephens". There have been only seven validated records of Brown Hawker, all since 1999 (and five in 2018-22). This species occurs close to the county boundary in Somerset and could be on the verge of colonizing Devon.

Warmer conditions in recent decades have enabled species to spread within Britain and to colonize from the Continent. Scarce Chaser *Libellula fulva* was first seen in Devon in 2003

and has since bred at several sites and wandered to many more, becoming especially widespread during hot conditions in 2022 and 2023. Small Red-eyed Damselfly *Erythromma viridulum* first appeared in Britain in 1999 and in south-east Devon in 2005, subsequently spreading as far as Plymouth. Red-veined Darters have increased, mainly as migrants but breeding has been recorded at Dawlish Warren (1995), South Huish Marsh (2003 and 2005) and Lower Bruckland Ponds (2006-7), while egg-laying was recorded in 2019 at both Clennon Valley Lakes and Meeth DWT Reserve and at Soar in 2022 and 2015; the species possibly bred at Blackhill Quarry (EDPH) in 2020. Lesser Emperor *Anax parthenope* has been recorded on at least 40 occasions since the first in 2003, with pairs egg-laying at Squabmoor Reservoir in 2006 and at Soar in 2022 (and in tandem in 2023). Vagrant Emperor has appeared on over 50 occasions and also more frequently in recent years, in part reflecting better awareness of rare dragonflies, especially among birdwatchers; in autumn 2013 a pair were seen at Bovey Heathfield and egg-laying was seen, only the second instance of this in Britain, and further egg-laying was seen at Soar in spring 2023 and East Prawle in autumn 2025.

In contrast to the significant northward range expansions that southern species have made within (and into) Britain in recent years, there is concern about the future of cold-adapted upland and boreal species like Common Hawker *Aeshna juncea* and Black Darter *Sympetrum danae*; such species may be lost from Devon as a result of climate change. These two species declined significantly in Britain during 1950-2019 (Taylor *et al.*, 2021) and are now included on the Red Lists for both Britain and Europe. It is noteworthy that thousands of small pools have been created in recent years in blanket bog in Dartmoor and Exmoor; initial results suggest that they are proving to be very attractive to Common Hawker and Black Darter, in particular, but also highlighting the presence of lowland species, such as Emperor Dragonfly, that may be breeding at higher altitude than formerly.

A Wandering Glider *Pantala flavescens* near Otterton in 2015, a Common Clubtail *Gomphus vulgatissimus* on the River Otter in 2017, a Norfolk Hawker *Aeshna isoceles* at Exminster Marshes in 2018 and a Southern Migrant Hawker at Beer in 2021 bring the total number of species recorded in the county to 40, of which 29-32, with the addition of Norfolk Hawker at Slapton Ley from 2022, are known to have bred since 2000.

## References

Cham, S., Nelson, B., Prentice, S., Smallshire, D. & Taylor, P. (eds.). 2014. *Atlas of Dragonflies in Britain and Ireland*. British Dragonfly Society & CEH Wallingford.

Merritt, R. 1983. An introduction to the study of dragonflies in Devon. *Nature in Devon* 4: 7-28.

Parfitt, E. 1879. The Fauna of Devon. *Transactions of the Devonshire Association*: 398-402.

Smallshire, D. 1996. An Inventory of Key Dragonfly Sites in Devon. British Dragonfly Society (Devon Group) – unpublished.

Taylor, P. *et al.* 2021. *Status of Dragonflies in Britain and Ireland 2021*. British Dragonfly Society, Old Weston, Huntingdon.

## Species Accounts

The species accounts below aim to summarise records up to the end of 2025, focussing on the main areas of occurrence and sites holding less common breeding species. Numbers are given (where the records contain these details) to indicate the relative abundance of species, but it should be remembered that these generally do not reflect the numbers that emerge from breeding sites; where useful data on productivity exist, exuvia counts are given. The sequence and species names used are those adopted by the British Dragonfly Society; alternative vernacular names are given in brackets.

### **Emerald Damselfly *Lestes sponsa* (Common Spreadwing)**

*Fairly common and widespread at standing waters*

Occurs at many sites with abundant rushes (*Juncus* species), including those at lower moorland elevations. National trends suggest that this is one of a small suite of species showing a decline over the 50 years from 1970; along with Common Hawker and Black Darter, Emerald Damselfly shows the greatest decline in site occupancy (a surrogate measure of population) in England, suggesting a common cause that is related to climate change. Counts exceeding 30 have come from Bovey Heathfield (50), Chudleigh Knighton Heath (40), Meeth Ball Clay Works (125), Petrockstowe Ball Clay Works (75), Smallhanger (100+), Stover CP (91) and Wixenford Quarry (Plymouth, 50), but none can match the estimated 800 at Combeshead Rewilding on 2 July 2023. Adults emerge typically between early June and mid-August (earliest on 23 May 2015 at Aylesbeare) and persist into September (latest on 2 October 2019 at Leftlake Mires).

### **Banded Demoiselle *Calopteryx splendens***

*Common and widespread at slow-flowing waters*

Found along the Rivers Axe, Carey, Clyst, Creedy, Culm, Deer, Exe, Lyner, Mere, Mole, Okement, Otter, Sid, Tamar (whole length), Taw, Torridge, Waldon, Weaver and Yeo. These include the lower sections of large rivers and, where conditions of suitable, the upper reaches and tributaries. Small numbers also occur in the inflows to both the Higher and Lower Leys at Slapton, believed to be the only breeding site in South Devon between the Exe and the Tamar. Often wanders away from its breeding sites to standing water such as at Dawlish Warren (five records, only one of them a male), Exeter Canal, Grand Western Canal, Lower Bruckland Ponds and Powderham Park, where breeding is unlikely. Absent from the Bovey Basin, although a male passed through Ventiford Ponds on 29 August 2021. Counts have included totals of about 100 or more in the lower Axe and Clyst valleys and Alphin Brook (Exeter), with over 500 along the Otter at Ottery St Mary. Flies from May (earliest 28 April 2008 near Hawkerland) to August, with a few into September (latest 11 October 1984 at Kilmington, River Axe).

## **Beautiful Demoiselle *Calopteryx virgo***

*Common and widespread, mainly at fast-flowing waters*

The highest counts – of 100 or more – have come from the Rivers Tamar and Torridge at Vealand and Volehouse Moor Nature Reserves, respectively, the fringes of Dartmoor (e.g. River Plym at Cadover Bridge, Meldon Reservoir, the Otter and Upper Teign Valleys, and the River Tavy) and the Rivers Otter and Teign. Like the previous species, individuals often wander away from breeding waters and are often seen flying over unsuitable standing waters; for example, there have been at least six records from Dawlish Warren and a single record from Lundy on 29 August 1985. Flies from late April (earliest 3 April in 1990, 2011 and 2016) to September (latest 10 October 2007 at Applethorn Slade).

## **White-legged Damselfly *Platycnemis pennipes* (Blue Featherleg)**

*Scarce along lowland rivers*

Typically found accompanying Banded Demoiselles, but more localised. Recorded from the catchments of the Rivers Axe, Carey, Clyst (including the leat at Clyst St Mary), Culm, Deer, Exe, Otter, Tamar, Taw, Torridge and Yeo. It occurs in slow-flowing sections with beds of emergent marginal plants and adults congregate in tall bankside herbage, such as nettlebeds. The highest counts have come from the lower Axe Valley, with 163 in 1995 and 255 (17 emerging) in 2007. Unusually, one has been seen emerging from the River Exe just below the tidal limit at Countess Wear. Like the demoiselles, this species may wander some distance from breeding sites, including to standing waters like Exeter Canal (where it is seen regularly and breeding is possible) and Lower Bruckland Ponds; in North Devon, 20 (3 pairs in cop.) were at Jennetts Reservoir in 2024 and over 100 were at Wooda Lakes, in the upper Tamar catchment near Dunsdon, in 2025.

Flies from late May (earliest 22 May 2007 at Exeter) to mid-August (latest 16 September 2007 at Cowley Bridge, River Exe).

## **Small Red Damselfly *Ceriagrion tenellum* (Small Red Damsel)**

*Scarce and local in acidic bog pools on moorland and heathland*

The main populations are on the fringes of Dartmoor and the East Devon Pebblebed Heaths, but the discovery of new sites suggests that this species is not as sedentary as formerly thought and dispersal may be assisted by bouts of increasingly high temperatures.

Known principally from the moorland fringes of Dartmoor, at Doetor Common (Doetor Mire/River Lyd/Walla Brook; maximum 80 in 1998), Gidleigh Common (Moortown Bottom; also a female in 2019 and seven in 2021 near Wallabrook Bridge, Scorhill Down, which is 370 metres ASL), Leftlake Clay Works (410 metres ASL – discovered in 2008, when six seen), near Piles Copse (discovered in 2011), Smallerhanger (near Cadover Bridge, discovered in 2018; 25 seen then and in 2024), the former stronghold at Smallhanger (the little remaining habitat there may be lost to clay-working in the near future), Wigford Down and Yellowmead (Willsworthy). Small but increasing numbers have been noted since at least 2022 at Emsworthy DWT, well away from the nearest known site, with about 15 (including

several mating pairs) in 2025. At least six, including mating, were discovered at Corndonford Bog in August 2025.

In East Devon, occurs at Bicton Common (maximum 34 in 2005, but only three in 2020) and nearby Blackhill Quarry, pools at Bystock Pools DWT Reserve (maxima 110 in 1994, 60 in 1995 and 36 in 2017) and Colaton Raleigh Common (has increased and spread in recent years, with maxima of 39 in 2020 and c50 in 2024). There were isolated records from Aylesbeare Common (in 1956, on 28 June 1965 and four on 7 July 2007) and then small numbers (maximum 19 in 2025) with breeding evidence in 2021-5. Also noted from Venn Ottery Common (1978); Woodbury Common (mating pair in 2009, probably from Colaton Raleigh Common), near Squabmoor Reservoir (a few in 2018, probably from Bystock Pools DWT Reserve) and at a garden pond near Otterton in 2024.

In North Devon, formerly known from Merton Moors (1978 and 1981); until recently, the only North Devon site was old settlement lagoons at Meeth DWT Reserve, where small numbers persist (14 in 2006, ten in 2008 and 25 in 2021), but two were seen at a new site in 2019 in at the DWT Beaver enclosure at Boldventure and in 2022 a large population was found nearby at Combeshead Rewilding, with at least 294 counted at six ponds; equally amazing was an estimated 580 at Combeshead in 2023. Also surprising was the presence of small numbers at Godborough, Bideford, since 2022; singles in 2023 elsewhere in NW Devon near Brandis Corner (a garden pond!) and the Hartland peninsula at East Titchberry, and recorded at Upcott Grange, Broadwoodwidger, in 2024.

Apart from Combeshead, the largest counts have come from Bystock Pools DWT Reserve (110 in 1994), Doetor Mire (80 in 1998); Moortown Bottom (200 in 1998 and 1999) and Smallhanger (170 in 1995 and 221 in 2005).

Occasional records from other Dartmoor sites have included: Burrator Arboretum (2002); Crownhill Down (2008, including a mating pair, but the site was subsequently lost to Hemerden Mine); Gutter Tor Mire (2002-3); Lee Moor (2005: two each at Big Pond and Whitehill Corner); Lydford Station (Railway Ponds – 20+ in 1993 and 1995, but no records since 2006; 70 were also found on the adjacent moor WSW of the old station in 1993); Narrator Brook (1997); Prewley Moor (small numbers in 2005-6, but not since); River Erme near Piles Copse (two in 2011); and Wigford Down (near Urgles: a few in 2006).

Old records come from the Bovey Basin at 'Bovey Tracey' (1921, 1925 and 1926); Stover (1950); Bovey Heathfield (1959); Chudleigh Knighton Heath (1965); Little Bradley Pond (1978-84, but the boggy pond was lost to natural succession); and a site 1.5km SSE of Chudleigh Knighton (1979 and 1981, but lost to Whitepit Quarry clay working in 1984).

Flies from June (earliest 1 June 2020 at Colaton Raleigh Common) to early September (latest 14 September 2002 at Bystock).

## **Southern Damselfly *Coenagrion mercuriale* (Mercury Bluet)**

*Rare: present at 7 sites in Dartmoor and EDPH*

In national and international terms, Southern Damselfly is the rarest dragonfly breeding in Devon. It and its habitats are protected by the Bonn Convention for the Conservation of Migratory Species of Wild Animals, the Berne Convention on the Conservation of European International Wildlife and Natural Habitats (1979), the EC Habitat and Species Directive (1992) and the Wildlife and Countryside Act (1981). It is also listed in the British Red Data Book and was a subject of the UK Biodiversity Action Plan (1994), which resulted in intensive research projects to determine the species' ecology and habitat management requirements. The Habitat and Species Directive has resulted in it being one of the qualifying features for inclusion in designated Special Areas of Conservation, which now cover the main breeding sites in Devon.

In Devon, it occurs at spring- and seepage-fed heathland streams and valley mires in water that is less acidic than normal at these locations, as it picks up dissolved base rich materials, is relatively warm and flows all year. Such sites are rare in Britain and typically very limited in extent, making the species highly vulnerable to reductions in habitat condition and hence local extinctions. Generic threats include heathland loss and fragmentation, undergazing and consequent scrub encroachment. One Dartmoor site (Tor View Moor) is on private enclosed farmland, while the other Devon sites are all in open heathland commons.



There are three extant colonies of Southern Damselfly on the East Devon Pebblebed Heaths. After records at Aylesbeare Common in 1956 and 1965, up to five were reported from a Black Bog-rush mire (NVC M14 *Schoenus nigricans* – *Narthecium ossifragum* mire community) in the RSPB reserve in most years during 1977-1992, apart from 13 in 1986. Thereafter, following a controlled burn, brush-cutting of Purple Moor-grass tussocks and the introduction of fenced cattle grazing, suitable habitat increased and peak transect counts rose dramatically to reach 165 by 2001. Following expansion to a newly suitable runnel in 2007 (burnt by RSPB three years earlier), the peak count rose to 204 in 2008 (34 in the extension area); numbers have been between 53 and 198 since, apart from a low of 15 in 2014.

It was first recorded at Colaton Raleigh Common in 1963, with up to 56 there during 1977-1989. Annual counts were made from 1991 and a formal transect established. Numbers peaked at 133 in 2005, but faltered after 2010, reducing to single figures since 2016, with only a single male in 2020, no more than two in 2021 and none since apart from one on 18 June 2025. This has been in spite of fenced cattle-gazing around the stream where the colony was originally focussed and the spring-fed flushes on the slope above where the damselflies spread to. There are concerns about the possible deterioration in micro-habitat quality on the slope in recent years, as well as concern over the reduction in stream flow following the repair of a leak in a pipe upstream that feeds water to Bicton Hall.

After records at Venn Ottery Common in 1979 and 1983, a colony there died out after the last was seen in 1990, apparently due to natural succession in the absence of grazing. Remedial management took place after the establishment of a Devon Wildlife Trust nature reserve, using techniques developed at Aylesbeare, and the UK's first licenced dragonfly re-establishment took place here in 2009 under the auspices of the Biodiversity Action Plan Steering Group and Liverpool University. This was originally planned for 2007, but due to a sudden reduction in water flow the release was aborted and 52 adults – 17 males and 35 females – brought from the Beaulieu area of the New Forest had to be released at Venn Ottery. Although the Aylesbeare Common colony lies just 2km to the south-west, natural re-colonisation from the increasing population there had not happened and studies elsewhere had shown that lifetime movement of mature adults was rarely over 500m. Studies had also shown that the genetic variability was weak in Devon, so it was decided to introduce more diversity from Hampshire, in the hope that the population might eventually mix with that at Aylesbeare. With better conditions in 2009, 500 adults were released in five batches, but it is notable that 23 adults also emerged from the 2007 release. Wisdom had it that the species matures in two years; indeed, none was seen in 2010 and the peak count in 2011 was 42, but two were seen in 2012, indicating that a few had matured in either one or three years. The species had clearly broken the biennial pattern, as transect counts during 2013-21 peaked at 98, 17, 98, 42, 62, 27, 39, 38 and 23, but only 14 in 2024.

Southern Damselfly also occurs at a few sites fringing Dartmoor, where it was discovered as recently as the 1990s following targeted searches funded by Dartmoor National Park Authority. DNPA has also funded systematic adult and larval counts in most years, the results of which have fed back into habitat management where necessary.

The first Dartmoor site was found in part of an extensive Rhôs pasture system at Tor View Moor, near Whiddon Down, in 1995. Peak transect counts ranged from eight to 41 during 1995-2004, then increased following management work by DNPA to peaks of 239 in 2008, 302 in 2015 and 422 in 2022. Although requiring DNPA effort to maintain good habitat condition at this small site, the management has resulted in good damselfly numbers.

Four km to the south-west of Tor View Moor, a colony was found in 1998 on the edge of Gidleigh Common, at Moortown Bottom. The damselflies were along three small watercourses flowing across an area of Gorse and grass where livestock had traditionally been fed in winter. With changes in livestock pressure, this site has subsequently needed considerable human input to replace the grazing and trampling effects of cattle, ponies and sheep. Peak transect counts of over 100 were made in 14 years during 1998-2024, and over 200 in eight years, with an all-time high of 292 in 2008. The northernmost runnel has tended to become overshaded, mainly by Purple Moor-grass, due to reduced trampling and grazing by livestock; this resulted in no larvae and only 3 adults being found in this runnel in 2024 and remedial action was planned.

Further targeted survey led to the discovery in 2005 of a colony over 11 km from Moortown Bottom, on the north-west fringe of Dartmoor at Prewley Moor. This easily accessible, sheep-grazed site has a series of small streams (NVC M29 runnels) flowing north-westwards from springs below a bracken-clad hillside. Uniquely for Devon, until recently this site also held small numbers of Scarce Blue-tailed and Small Red Damselflies. Peak transect counts have exceeded 100 each year (except for 44 in 2016), with an impressive high of 386 in the

fine summer of 2023. Some dredging work was implemented in August 2016 to remove silt and vegetation and improve the conditions in a drain that intercepts most of the hillside streams (unfortunately, this led to the loss of adjacent boggy areas that hosted the Scarce Blue-tailed and Small Red Damselflies).

A scrubby extension of Prewley Moor across the A386 road at Lower Prewley has held small numbers since 2008, with peak counts of 36 in 2009 and 41 in 2021. Presumably resulting from wandering individuals from the Prewley Moor colony some 850m away, it benefitted in 2008/9 from scrub management to maintain an open channel for the species, which is a critical element of the habitat condition.

In the Blackdown Hills, Southern Damselfly was recorded in springline mires at Southey Moor (Smeathorpe), just across the border into Somerset from Gotleigh Moor, on 14 July 1959 and at Hense Moor on 29 June 1965. However, subsequent searches failed to find the species, which was probably lost as a result of a lack of grazing and/or scrub encroachment of the boggy runnels (the fate of many springline mires in the Blackdowns). Following eight years of scrub control and grazing management under Higher Level Stewardship, Hense Moor was assessed early in 2015 as being suitable for Southern Damselfly re-establishment and under licence, 460 were transferred from the New Forest in June of that year. Unfortunately, this second translocation targeted under the UK Biodiversity Action Plan failed, presumably due to unsuitable or insufficient micro-habitat to support the species.

Flies from late May (earliest 25 May 2010 and 2020 at Colaton Raleigh Common) to mid-August (latest 3 September 1998 at Moortown Bottom).

### **Azure Damselfly *Coenagrion puella* (Azure Bluet)**

*Very common and widespread*

Most abundant in well-vegetated ditches, ponds and the edges of lakes, where egg-laying tandem pairs can be seen in large numbers. Counts at Stover CP have exceeded 1000 on occasions, with over 1000 estimated on 2 June 2020 (about 800 of these were attracted to the luxuriant growth of pondweed in the lake) and 2180 in 2023. About 1000 were estimated to be at Exminster Marshes on 7 July 2019. Flies from late April (earliest 5 April 2016 at Prawle) to August, with a few into September (latest 19 October at Otterton).

### **[Variable Damselfly *Coenagrion pulchellum* (Variable Bluet)]**

*Status uncertain*

A putative blue-form female was photographed at the western fringe of Braunton Marsh on 19 May 2023. The species occurs in the grazing marsh ditches of the Somerset Levels but its presence in Devon has never been verified, despite records in the Transactions of the Devonshire Association from 1879 and 1976-80. Females can be easily confused with Azure Damselfly, but the Braunton Marsh image (right) appears to show a tri-lobed rear margin to the pronotum, although lacks a pale bar between the eyes.]



## **Common Blue Damselfly *Enallagma cyathigerum* (Common Bluet)**

*Very common and widespread*

Most abundant at large ponds, lakes, reservoirs and canals, but sometimes on slow-flowing waters. The species disperses widely, though, and may turn up anywhere. Males of this species are the most likely blue damselflies to be seen well out over water, perched on surface vegetation or patrolling close to the surface. Peak numbers are typically later in the summer than Azure Damselfly, with transect counts totalling 400+ in 2010 at Stover CP and 785 estimated there on 12 August 2020. Counts of more than 500 have come from Exeter Canal, Grand Western Canal, Lower Bruckland Ponds, Meeth Quarry, Rackerhayes and Seaton Marshes. Up to 200 have been recorded on Lundy, where the species has bred. Flies from May (earliest 20 April 2015 at Lower Bruckland Ponds) to September (latest 17 October 2021 at Bovey Heath).

## **Red-eyed Damselfly *Erythromma najas* (Large Redeye)**

*Fairly common but very localised at standing waters with floating vegetation*

Known mainly from the Bovey Basin, the lower Exe Valley and Grand Western Canal, especially where there are water-lilies, on which the males sit in strategic positions. Has expanded its range considerably in recent decades.

In the Bovey Basin, large numbers have been counted at fishing ponds at Rackerhayes, Kingsteignton, since 1978, with 'hundreds' in 1989, 220+ in 1995, 173 in 2005 and 516 on 13 June 2021, but very few in 2025 following dredging of the lake. The main site with public access is Stover CP, where over 100 (1995 and 2006) were noted on the lake until the loss of water-lilies around 2009; no more than six were seen during 2011-17, when it was mainly restricted to a small area of non-native water-lily at the edge of the lake and in surrounding ponds; perhaps resulting from an improvement in water quality, there was a rapid growth of pondweed in the lake after 2018, resulting in counts of 119 in 2020 and 436 on 14 June 2021. In addition, small numbers breed at Abbrook Pond (21+ in 1992), in Decoy CP at Decoy Lake (100+ in 1992, since when a maximum of 61 in 2025) and Magazine Pond (20 in 1992 and 2023), Eddison Pond (20+ in 1992, but no more than five since), Gallows Cross SUDS ponds, Kingsteignton (20 in 2020, 12 in 2021, but none since 2022), Little Bradley Ponds (1978, up to five 1992-2021, with egg-laying in 2018, rising to 22 in 2022 and 43, with six pairs egg-laying, in 2023 and 45 in 2024), Netherexe Pond (at least 30 in 1995 and 2006; also present at the former pond at this site, known as Key Transport, in 1978 and 1986) and Ventiford Ponds (private ponds close to Stover created at a site previously known as Alder Park, where 20 were recorded in 1995; noted at the new ponds from 2013, with maxima of 49 in 2020, 93 in 2021 and 125 on 27 May 2022). Occasional records have come from Bovey Heath (2009, 2022), Bradley Pond (1987), East Golds (Kingsteignton, 2005 and 2021), Sawmills Pond (1990-2005) and West Golds (1992). It was formerly present (1982) at the Centrax Pool, Newton Abbot, until the site was lost to development.

In the lower Exe Valley, Red-eyed Damselfly is found along the whole length of Exeter Canal, with 100+ near Countess Wear Sewage Works alone in 2009 and 320 in 2023; its presence has continued despite the control of Fringed Water-lily and regular pondweed cutting to

maintain a boat channel. Small numbers, no doubt originating from the canal, have been found on the wider ditches of Exminster Marshes in recent years, and 16 were found on the lagoon there in 2006. Small numbers have also been seen on the River Exe, Harpers Fishing Ponds (Double Locks), Matford RSPB pools (2005), Dawlish Warren (2013 and 2023-25, with at least 10 emerging in 2024), Hazelwood Holiday Park (2023), Topsham (2017) and Dart's Farm (since 2022, with 20 in 2024).

It was first recorded along the Grand Western Canal in 2004 (20), with transect counts along the whole length totalling 118 in 2006, 502 in 2010 and 215 in just two sections in 2018. There are isolated breeding colonies in East Devon at Squabmoor Reservoir (up to 20 since 2005) and Lower Bruckland Ponds (maximum 33 in 2008), while a single male was at Blackhill Quarry in 2020 and two were at Bystock in 2023 and small numbers in 2024.

In North Devon, a small breeding population was reported near Beaford in 2008 and at least six were at Lake, near Shebbear, in 2022; both sites are some 50km W or NW of the nearest known sites. In 2024-25, 2-5 were at Wooda Lakes, near Dunsdon, in May-June, while a similar number were at Gammaton on 4 July 2025.

Flies from May (earliest 3 May 2001 at Exeter Canal) to August (latest 18 September 1982 at Rackerhayes).

### **Small Red-eyed Damselfly *Erythromma viridulum* (Small Redeye)**

*Recent colonist, breeding at standing waters across southern Devon*

This species has shown a dramatic expansion in range in Britain since it was first recorded in Essex in 1999. It soon reached south-west England, with the first in Devon in 2006 (maximum 54) at Lower Bruckland Ponds, where there were 105 in 2008 and at least 310 on 20 July 2018. Next it appeared at Exeter Canal in 2009 (15, with 25 in 2014, 35 in the section by Countess Wear Sewage Works in 2019, and over 600 on 17 July 2022), Dawlish Warren in 2010 (maximum of 20 in 2014), Seaton Wetlands in 2013 and at many more location since.

The first records west of Exeter Canal came in 2017, with records from three Bovey Basin sites, including 42 at Little Bradley Ponds (with 50+ there in 2018, 61 in 2020) and eight at Ventiford Ponds (with 27 there in 2018 and 71 in 2022). Noted in 2018 at four other Bovey Basin sites, including the first records for Stover CP (seven in August 2018, rising rapidly to 100 in 2019 and 577 in 2022). Surveys at Rackerhayes found 311 on 24 August 2021.

In East Devon, 100+ appeared at Squabmoor Reservoir on 22 July 2018, but only ten were present three days later, suggesting that most had moved on quickly. Well outside the established range, it was first found in 2018 at Shobrooke Park fishing lakes (30+) and Clennon Valley Lakes (40+) and in 2019 at Drakes Reservoir, central Plymouth (several). In 2020, three were on the coast at Starehole Ponds, Soar, from 11 August to 1 September; in 2021 small numbers were noted at new sites at Bishopsteignton and Bolberry; and in 2022 there were 30+ at Gammaton Reservoirs, Bideford, 19 at Southdown Valley, Soar, and singles at Aylesbeare Common, Langtree and Slapton Ley. In 2023, also noted at Grand Western Canal and Lower Otter Restoration Project; in 2024-25 at Darracott and Gammaton (up to 10) Reservoirs, near Great Torrington; and in 2025 at Godborough, Bideford.

Flies from late June (earliest 22 June 2023 at Ventiford Ponds) to early September (latest 16 September 2007 at Lower Bruckland Ponds).

It is very likely that other breeding sites exist across the county, given the frequent occurrence of suitable ponds with floating pondweed or algae, and further spread seems inevitable. However, the species is easily overlooked and the use of binoculars is almost essential to detect and identify this and the preceding species when they are out over the water.

### **Blue-tailed Damselfly *Ischnura elegans* (Common Bluetail)**

*Common and widespread in standing and slow-flowing waters*

This, one of the most widely recorded damselflies in Britain, has a low habitat tolerance threshold and hence tends to be commonest at wetlands with relatively poor water quality. It is generally scarcer at the higher quality sites, but maximum recorded at Stover CP is 398 in 2025. Estimates of over 500 come from Seaton Marshes in 2001, Slapton Ley in 2006 and Ventiford Pools in 2018. Up to 20 adults have been seen on Lundy, where the species has bred. Flies from late April (earliest 20 April at Slapton Ley in 2011 and Braunton Burrows in 2019) to early September (latest 11 October 1978 near Sparkwell).

There may be a link between a significant decline in site occupancy in England and the recent widespread use in arable crops of neonicotinoids, which have been shown experimentally to reduce the productivity of this species.

### **Scarce Blue-tailed Damselfly *Ischnura pumilio* (Small Bluetail)**

*Scarce and local*

This Red Listed specialist of early successional stages has nationally important populations in Devon. Its often ephemeral occurrences and a need for sparse vegetation cover make its long-term conservation rather difficult. Historically, it was known principally from both working and former clay pits in the Bovey Basin, Lee Moor and Meeth-Petrockstowe areas, where its current status is uncertain but certainly less favourable than previously. Scarce Blue-tailed Damselfly was discovered during the 1990s at a number of muddy streams and ponds in Dartmoor moorland, especially where livestock trampled ('poached') bankside vegetation at crossing points; with reductions in stocking densities, especially in winter, such sites may now be less suitable and some have likely lost the species. By 2025, there were some 18 sites where no records had been received during the previous 10-20 years.

Dartmoor breeding sites include the following:

Cadover Bridge-Blackaton Cross area: the old Brisworthy clay settlement lagoon at Cadover Bridge (first recorded in 1978; maxima 45 in 1993, 77 in 2005, 51 marked in 2006, 25 in 2010 and 2018) and Blacka Brook (maximum 16 in 2005) up to Blackaton Cross (35+ in 2018); presumed wanderers (four) from Cadover were found nearby at Shaugh Moor 2019 and at least four males were found at the adjacent Smallerhanger in 2024.

Corndonford Bog: up to five during 1995-2006.

Doetor Common: small numbers during 1995-98.

Gutter Tor Mire-Sheepstor: discovered in 1995, maximum 40 in 1999, but not recorded since.

Lee Moor claypits complex: Cholwich Town Waste, Quick Settling Ponds and Tory Brook in 1996-7, Whitehill Corner in 1996 and 2005, Portworthy and Refinery Settling Pond in 2005, with over 100 at the latter; these records pre-date the cessation of mining and pumping and the species may no longer occur here, although it persists in the contiguous Cadover Bridge-Blackaton Cross area.

Legis Mire: small numbers in 1995, 1998 and 2005.

Lud Brook Head, Ugborough Beacon: recorded in 1995, 1998 and 2005.

Lydford Station old railway ponds: 'very good numbers' recorded in the late 1970's-early 1980's, 30 in 1988-89, only six by 1993, when the favoured area had become overgrown, and only four in 1995; seven caught and marked in 2006 is the only subsequent record.

Prewley Moor: small numbers during 2005-15, with at least six in 2009 and four males marked in 2006.

Simon's Lake, Sherberton Common: small numbers in 1995 and 1997.

Small Brook, near Avon Dam: up to three in 1995 and 2005.

Smallhanger Waste: the first records at this former stronghold were during 1978-80; counts include 54 in 1995; 20 in 2001; a maximum of only seven during 2002-5; 30 in 2006, including 29 marked in a study; 20 in 2010; and no more than ten subsequently as the reworked area has expanded.

Walla Brook, Riddon Ridge: recorded during 1995-2006, with 95 in 1999.

Whitchurch Down: small numbers during 1986-98.

Wigford Down (streams on NW fringes): recorded occasionally since 1995, with a maximum of 10 in 1998 and 2018.

Vixen Tor Mire: small numbers in 1995 and 1998.

Isolated Dartmoor records have come from Burrator Arboretum (2004), Challacombe Farm (2006), Cornwood (just off the moor, 1978), Mary Tavy Reservoir (just off the moor, 1978) and Meldon Aplite Quarry (2018-9, egg-laying seen in 2019).

In the Bovey Basin, permitted surveys have resulted in the discovery of two colonies in working ball clay pits. At Chudleigh Knighton Clay Pit, 80 were seen in 1994 and 31 in 1996, when reworking destroyed the breeding area; 16 (11 males) were translocated to Little Bradley Ponds on 6-7 July 1996, but none was seen subsequently (although there were records there in 1999). At East Golds, up to six were seen in 1995-6 and two males in 2021. There have also been a few records of small numbers from the fringes of other clay pits, including Whitepit Quarry (three in 2006), as well as at nearby ponds at Chudleigh Knighton Heath (two immature males in 2006), Finlake (two immature males in 2009), Bovey Heathfield (one or two at temporary puddles in 2005-6) and Little Bradley Ponds (in 1980, 1982 and two males on 21 August and a female on 28 August 1999). An old record from Stover in 1970 indicates the early presence of this species in the Bovey Basin.

Following a record from Orchard Meadows, Saltram, Plymouth in 1996, there were records during 2001-7 from a colony almost at sea level nearby at Wixenford Quarry. There were 18 here on 30 July 2007, including several fresh, immature specimens and two exuviae, suggesting a second generation.

In North Devon, it has been recorded from old ball clay workings at Meeth Quarry (68 in 2006 and 23 in 2008 in systematic surveys, and last seen in 2016; now flooded and a DWT Reserve) and nearby at Petrockstowe Ball Clay Works, where nine were seen in 2005, but it is doubtful if the species persists at either site. Isolated records have come from Bridgerule Mill in 1990 and 1992 and ponds at Welcome (up to seven in 1992 and 1998, with breeding proven) and Heddon Valley, Exmoor, during 2009-11, nearly 40 km NE of the nearest known site.

Flies from June (earliest 19 May 2011 at Heddon Valley) to August. The latest records have both been on 31 August and are suggestive of a second generation: a well-documented female in 2012 at Tor Woods, Salcombe (perhaps of Continental origin and coinciding with records on the Norfolk coast around the same time), and in 1996 at Saltram (near to which a possible second generation was recorded at Wixenford).

### **Large Red Damselfly *Pyrrhosoma nymphula* (Large Red Damsel)**

*Common and widespread in standing and slow-flowing waters*

Typically the first species to emerge in lowland areas in spring, this species also breeds commonly in ponds and both blanket and valley bogs in moorland, where the flight season is delayed. Over 500 were estimated at Vixen Tor Mire on 9 July 2005. Maximum counts at Stover CP of 200+ in 2000, 100+ in 2018 and 106 in 2020. There have been counts of over 100 at Bicton Common, Bystock DWT Reserve, Castor Down, Colaton Raleigh Common, Heddon Valley, Lee Moor, Little Bradley Ponds, Meeth Quarry, New Cross Pond, Ottery St Mary, Rackerhayes, Slapton Ley, Smallhanger and Vealand.

Flies from April (earliest 31 March at Orley Common) to early September (latest 23 September 1978 at Bystock).

### **Southern Migrant Hawker *Aeshna affinis* (Blue-eyed Hawker)**

*Vagrant: two records*

A photograph taken of a very blue hawker at East Soar on 6 September 2018 wasn't quite good enough to enable this species to be added to the Devon list, but the same cannot be said of a male photographed at Beer Head on 2 August 2021 (image: Steve Waite). A male was seen at Southdown Valley, Soar, on 21 August 2025.



## **Southern Hawker *Aeshna cyanea* (Blue Hawker)**

*Common at lowland standing waters, including garden and woodland ponds*

Southern Hawkers are widely distributed across Devon, including the moorland fringes. This large, brightly-coloured dragonfly often attracts attention when a territorial male 'challenges' a human intruder, hovering close by. Females may be watched at close quarters egg-laying at the edges of ponds. Although most adults are seen at breeding sites from July until October or November, they emerge during late May-July and move to woodland to feed and mature and are often seen hawking in woodland rides and glades. Emerges from June (earliest on 23 May 2009 at Stover), flying at territories mainly from July to October, with some into November (latest 12 December 2006 at Ernesettle). On Lundy, one was seen on 30 August 1999 and two during 12 August-26 September.

## **Brown Hawker *Aeshna grandis***

*Rare visitor: nine records*

Although Brown Hawker is quite common just over the border in Somerset and Dorset, its rarity in Devon (and Cornwall) is an enigma. Singles were at Little Bradley Ponds on 26-28 August 1999; Ideford Common on 3 September 2012; the Grand Western Canal on 6 July 2018 and 3 September 2021 (the canal is a good route for colonisation from known sites a few kilometres over the border near Taunton, Somerset); in a garden at Manaton in 2019 (probably in August); at Roborough (near Torrington) on 11 July 2020; in a garden at West Hill on 1 August 2021; at Godbrough, Bideford, on 5 July 2022; and at Yonder Oak Wood on 30 August 2024 (awaiting confirmation).

## **Common Hawker *Aeshna juncea* (Moorland Hawker)**

*Fairly common on moorland, scarce (or absent?) in lowland heathland*

Records of this species are marred by confusion with the very similar and much more widespread Migrant Hawker, although this is a large and powerful species and clearly does wander widely, especially across its preferred moorland habitat. Most of the confirmed Common Hawker records come from the uplands of Dartmoor and, to a lesser extent, Exmoor, where the species inhabits acidic ponds and open water in valley mires and blanket bogs. It is also known from ponds and boggy areas remaining from old clay and tin workings around Dartmoor. However, records away from the moors have dwindled in recent years, to the extent that the species may no longer breed below about 250 m ASL. National trends suggest that this is one of a small suite of species showing a decline 1970-2019; along with Emerald Damselfly and Black Darter, Common Hawker has declined in site occupancy (a surrogate for population) in Britain, implying a cause related to climate change (Taylor *et al.*, 2021).

Dartmoor blanket bog sites include: Cowsic Head (1978); Cramber Pool (2010); Plym Head (1984); Quickbeam Hill (Petre's Pit, 1986 and 2020); Skir Hill (1986); and Taw Marsh (1978 and 1996). Recent peatland restoration work, involving the creation of many hundreds of pools in the Okement Hill-Great Kneeset-Cranmere Pool area has started to boost numbers of this species (and Black Darter). Records from these remote, high altitude areas are

sparse, due to access difficulties. There are more sightings from the more accessible lower mires and pools, mainly around the moorland fringe. Other main sites include: Blackabrook-Blackaton Cross-Lee Moor area (maximum 16+ in 2005, breeding in 2010, still three in 2018); Blackaton Brook (Raybarrow Pool-Shilley Pool-Kennon Hill, 1977, 1986, 2011); Burrator Arboretum (1977, 1987, 2002, 2005); Cadover Bridge area (1978-2022, breeding in 1995, 2018 and 2022); Gidleigh Common (Moortown Bottom and Walla Brook, 1996-2005); Haytor Quarry (1978-2004, but none reported since since); Leftlake Mires (up to six in 1985 and 2008-19); Lydford Railway Ponds (1978-2001, with egg-laying in the last year); Raybarrow Pool-Kennon Hill (1977, 1986); Red Lake China Clay Works (1977-2018); Small Brook (near Avon Dam, 1995-2016, with egg-laying in 1995); Smallhanger (1978-2012, with up to 40 recorded and breeding evidence in eight years, but the site has been largely lost to reworking since); and Statt's Bridge (1996, 2012-25, mating in 2016, egglaying in 2025).

Less frequently recorded Dartmoor sites include: Aish Tor (2025), Believer (1983), Blackslade Mire (2016), Boldventure (near Chagford Common, 2003), Chagford (1950), Coombe Down (1978 and 2017), Cramber Pool (2019), Crownhill Down (breeding in 2008, site destroyed by Hemerdon Mine), Cordonford (2002), Dart Valley (Hannaford Manor, 1982), Emsworthy DWT Reserve (2016-17, egg-laying in 2021-25), Fernworthy area (1950, 1981, 1983, 2025), Foggintor Quarry (1982), Hanger Down (1978), Harford (Tor Rocks Quarry, 1981 and 1988), Headon Down (2011), Holming Beam (2019), Houndtor (Hedge Barton, 1978), Huntingdon Warren (1979), Lee Moor (1989), Lower Merripit (2022), Lud Brook Head (2005), Meldon Reservoir (Vellake Corner, 2003), Merrivale Quarry (1982), Newbridge Marsh (2002), New Waste (1988), Nun's Cross (2016), O Brook (2015), Penn Moor (2024), Pixie's Garden Common (Ramsley, 2005), near Postbridge (2012), Prewley Moor (2005, 2012), Quick Settling Ponds (Lee Moor, 1997), Rattle Brook Peatworks (2019), River Lyd (High Down, 1995), Swincombe Valley (1978), Upper Erne Valley (2009, 2011), Thornworthy (1981), Vogwell (2012), the Walkham Valley (Bedford Bridge 1983, Grenofen 1978 and Heckwood Quarry 1980-1), West Blackdown (1981, 1990 and 1995) and West Glaze Brook (2016). Perhaps a wanderer from Dartmoor, a male was photographed in 2012 at Grenoven Wood in the lower Tamar Valley.

There are many fewer records from Exmoor than from Dartmoor, probably due to the poorer condition of the blanket bog – although there have been efforts to restore bogs in recent years as on Dartmoor. Exmoor reports come from Bray Reservoir (1992), Challacombe Common (1999), Farley Hill and Farley Water (1982), Heddon Valley (2009) and Litton Water (1968), while the species occurs just over the Somerset border in the Pinkery Pond and Squallacombe areas.

Other records from North Devon have come from Braunton Burrows (1979), Brendan and Vealand Fen (an emergent in 2009), Bude Aqueduct (1995), Coombe Mill Pond (2007), Meeth Quarry (2010-11), Petrockstowe Ball Clay Works (Courtmoor Quarry lagoons, 2005), Sterridge Valley (2009), Witheridge Moor (2006-23) and Winkleigh (Wheatland Farm, 2009). There is an isolated record from the Grand Western Canal (2005).

Ponds in the Bovey Basin heaths are generally not acidic, due to the neutralising effect of the underlying ball clay deposits, but nevertheless Common Hawker may still have a tenuous foothold here. There are breeding records from Bovey Heathfield (2009-10, with others seen in 1988 and 2008-20), Chudleigh Knighton Heath (1993-2012, but not since) and at Stover CP (records in 1978, 1982, 1985, 2002, 2007, a territorial male in 2009 and a mating pair in 2010). Nearby, isolated reports have come from Bradley Manor (Newton Abbot, 1985), the old Key Transport Pond (now Netherexe Pond, 1978), Little Bradley Ponds (1978, 1992 and 2012) and Rackerhayes (1975 and 1985). There were records from Haldon Forest during 1979-95, with evidence of breeding at two different sites in 1989 and 1995, but no records since then. Surprisingly, there are no breeding records from heathland ponds in the EDPH, although there are reports from Aylesbeare Common (in 1977, 1994, 1996, 2010, 2011, 2013-4 and 2016), Bicton Common (1977, 1981 and 2019), Venn Ottery Common (1979, 1983 and 1987) and Woodbury Common (2010).

Reports from the south Devon coastal fringe, at Seaton Marshes, Slapton Ley, Wembury and between Exeter and Dawlish Warren, must carry elements of doubt, due to the abundance of Migrant Hawkers in these areas, although one was confirmed at Dawlish Warren in 1994. The same might apply to records of three singles on Lundy in September-October 2002-28.

Emerges from late June (earliest 4 June 2011 at Witheridge Moor), flying until September (latest 17 October 2021 at Bovey Heath).

### **Migrant Hawker *Aeshna mixta***

*Fairly common lowland breeder and migrant*

Prior to 1990, Migrant Hawkers were largely restricted to sites within 15 km of the south coast. Since then records have come from most parts of the county, although only from the lowest fringes of Dartmoor (e.g. Smallhanger). Tolerating waters with elevated conductivity and salinity levels, the species can be abundant amongst emergent vegetation around ponds, lakes and canals, and in reedy ditch systems in coastal marshes. Counts of over 20 have come from Goosemoor, Otter Marshes, Slapton Ley, South Huish Marsh and Stover CP, with maxima of 55 at Grand Western Canal (2005), 60 at Exminster Marshes (2005) and 73 at Seaton Wetlands (2001). Over 20 exuviae were found at Dawlish Warren on 10 August 1992 and 55 at Stover CP on 18 August 2024. Presumably still also occurring as a migrant, numbers can often be seen at coastal locations away from water, sometimes flying over the sea. Up to three have been seen on Lundy. Flies from mid-late July (earliest 30 June 1995 at Cadover Bridge) to early November (latest 18 November at Broadsands in 2005 and Dawlish Warren in 2006).

### **Norfolk Hawker *Isoaeschna isoceles* (Green-eyed Hawker)**

*Rare visitor and presumed breeder*

One frequented reedy drainage ditches at Exminster Marshes on 12-13 July 2018 (pictured). This species has expanded from its traditional East Anglian haunts in the last decade or so to North Kent, the East Midlands and RSPB Radipole Lake, Dorset. Sightings came from the Avalon Marshes, Somerset, and coincidentally in south Devon from Slapton Ley in 2022.

Up to nine were seen at Slapton Ley between 17 June and 29 July 2022, including mating pairs and egg-laying females on at least three occasions, with sightings concentrated in the area around Slapton Bridge, mostly around the dipping platform; up to nine were seen on 27 June. Given the multiple occurrences, it is possible that the species may have arrived and bred two years earlier. In 2023, singles were seen near Slapton Bridge on 28 May and at the southern end of the Lower Ley in a garden at Torcross on the 30<sup>th</sup>, followed by five near the Slapton Bridge on 2 June and about 12 in the same area on 13 June. In 2024, individuals were near the southern end of Slapton Ley at Stokeley Farm (filmed) in June and three at Beesands Ley on 7 June, suggesting further spread. Up to three were at Slapton Ley in June-July, with a late individual up to 16 August. In 2025, five males were at Slapton on 20 June, with at least four still flying on 4 July.



In south-east Devon, singles were at Lower Bruckland Ponds on 16-17 June 2023, 21 July 2024 and 4 July 2025, while one was recorded at Exminster Marshes during 26-30 June 2025.

### **Vagrant Emperor *Anax ephippiger***

*Rare vagrant; has attempted to breed*

In keeping with most UK records, this wanderer from southern Europe or (more likely) North Africa has appeared on over 50 occasions, during 21 February-15 June and 9 September-24 December, all but two since 2011. Reports often coincide with warm, southerly airflows and sometimes 'Saharan' dust. Records (singles unless stated otherwise), in chronological sequence, are as follows:

1903: Devonport on 24 February, the first documented record.

1988: photographed in a garden at East Prawle on 8 November.

2011: found hanging on washing in a garden in Oreston, Plymouth, on 15 April, heralding an unprecedented spring influx in southern Britain; a male Lesser or Vagrant Emperor was over Exeter Canal by Exminster Marshes on 2 October.

2013: at and around a pond at Bovey Heath, two were first seen on 8 October, at least one of which was a male; a male remained to 2 November; a female was seen egg-laying on 18 and 26 October (image: Pauline Smale), the first such record in Britain since one in 2011 in Cornwall;



and a female last seen on 12 November. Despite checks the following year, only Emperor Dragonflies were found to emerge.

2015: a male on Lundy on 13 October and a female found in an airing cupboard at Roborough, near Torrington, also on 13 October, presumably having been brought in with washing the previous day; one at Coypool, Plymouth, on 5 November.

2017: a female or immature (presumed to be of this species) was at Mansands Beach on 8 April and singles at Dawlish Warren on 13 May and 8 October.

2018: one at Dawlish Warren on 18 October.

2019: singles at Cornwood (presumed to be this species) on 21 February; Plymouth and Lundy on 25 February; between Stoke Point and Mothecombe Beach on 26 February; Soar Mill Cove on 27 February; Budleigh Salterton (presumed) on 11 March; Mutley Plain (presumed) on 18 March; in a garden at Cockington, Torquay, (presumed) on 25 March; Oreston on 26 March; a female at Dawlish Warren on 16 October; and a male photographed at Gammaton Moor on 27 October.

2020: a male was found on Lundy on 11 October and a moribund individual was found on the edge of Bideford on 5 November.

2021: singles at Prawle Point (caught in a moth trap) on the night of 9-10 September, Berry Head on 11 September and 30 October, Exminster Marshes on 8 October, Dawlish Warren on 3 November, Stoke Point on 7 November and Lipson, Plymouth, on 13 November.

2022: one at Preston Park on 10 September and a female at Bovey Heath on 17 September.

2023: a female or immature (possibly 2?) at Dawlish Warren on 10 April, a male in the dunes there on 13 April, a probable female on 10 May, a male and female on 11<sup>th</sup>, two males on the 13<sup>th</sup> and one male on the 20<sup>th</sup>; a female or immature at Slapton Ley on 23-25 April; single males were seen at Wembury on 12 April and 12 May, near Otterton on 20 May and at Bowling Green Marsh and Mansands on 21 May; an exceptional five were at Southdown Valley Lagoons, Soar, on 24 May, comprising three males and a tandem pair ovipositing, with one male remaining on the 25<sup>th</sup> and 4 June; coincidentally, at least four were on Lundy on 24 May with at least two males remaining on the 25-26<sup>th</sup>. Later in the year, singles were at University of Plymouth on 12 October; Sticklepath, Barnstaple, on 17 October (dead); Lundy on 19 October; and at Sidmouth Croquet Club on 19 November.

2024: a male in Plymouth on 18 December and individuals presumed to be of this species near Chudleigh and at Dawlish Warren on 24 December.

2025: a male at Southdown Valley Lagoons on 20 & 22 May; two at Beesands Ley on 15 June; a pair was seen mating and egg-laying at Moorwell Pond, East Prawle on 9 October (image: Pat Mayer); singles at Dawlish Warren on 14 October and 6 and 8 November; on last date also singles at Teignmouth/Shaldon (location to be confirmed) and (female) at Berry Head.



In addition, there have been occasional reports of dragonflies likely to have been this species, especially those 'out of season'. Individuals were seen in the dunes at Dawlish Warren on 28 April and 16 May 2011, Paignton on 12 November 2013, Exeter on 7 April 2017, Dawlish Warren on 3 November 2019 and The Barbican, Plymouth, on 11 January 2023.

### **Emperor Dragonfly *Anax imperator* (Blue Emperor)**

*Common resident; breeds widely in standing waters*

Widespread across the county, there have been range extensions to some upland Dartmoor and Exmoor sites in recent decades, in line with very significant increases in Britain over the last 50 years. Although this species is highly territorial at breeding sites, transects and other systematic counts at sites with series of ponds or ditch systems have produced totals of up to 25 at Exminster Marshes (2006), 21 (2010) along the Grand Western Canal, 30 at Lower Bruckland Ponds (2006), 17 at Rackerhayes (2005), 20 (1998, 2000 and 2009) at Smallhanger and 19 (in 2020) at Stover CP. Seen occasionally on Lundy in July-September. Counts of exuviae include 41 from an Exeter garden pond in May 2020 and 54 at Ventiford Pools on 22 June 2023.

Flies from mid-May (earliest 5 May 2020 at Dawlish Warren) to September (latest 29 October 2025 at Umboorne). There has been an increase in late-season records in recent years, with all 14 of the October records since 1993.

### **Lesser Emperor *Anax parthenope***

*Rare migrant; has attempted to breed*

There have been over 40 records in Devon of this recent British colonist in 15 years since the first in 2003, with egg-laying in 2005, 2022 and 2023 and most records in the hot summers of 2006, 2019 and 2023. Records have been between 11 June and 18 September; therefore a male Lesser or Vagrant Emperor over Exeter Canal by Exminster Marshes on 2 October 2011 is more likely on timing to have been the latter. Unless specified otherwise, all have been single males, which are typically driven away from potential territories by male Emperor Dragonflies:

2003: two at Smallhanger on 10 August.

2005: Exeter Canal at Countess Wear on 22-23 June; a pair egg-laying (the male was taken by a Moorhen and fed to a chick!) at Squabmoor Reservoir on 17 July.

2006: Exminster Marshes on 16 June; Little Bradley Ponds on 17 June; Squabmoor Reservoir during 24-30 June; Lower Bruckland Ponds during 16-21 July; and Okenbury, near Kingston, on 9 August.

2010: Aylesbeare Common on 8 August and Smallhanger on 30 August (two male, one female).

2012: Lower Bruckland Ponds on 10 August.

2014: Bowling Green Marsh on 1 July.

2017: Mill Street, Ottery St Mary on 18 July.

2018: Netherexe Pond on 22 July.

2019: Lower Bruckland Ponds on 28 June and 2 July; Stover CP on 6 July, 15 July and 2 August; Slapton Ley and Squabmoor Reservoir (or possibly Vagrant Emperor) on 10 July; Meeth DWT Reserve on 14 July and a female photographed at Croyde Beach on 24 August.

2020: Dawlish Warren on 3 August female); Stover CP on 11-13 August (two separate males, one pictured); and Venford Reservoir (Dartmoor) on 12-13 August.

2021: Lower Bruckland Ponds on 21-22 July, Slapton Ley on 2 August and Meeth on 4 August.

2022: Grand Western Canal on 11 June and Rewilding Roborough Fields on 12 July, while at Southdown Valley, Soar, a male on 29 July and 27 August and an ovipositing pair in tandem on 8 August.

2023: a good series at Southdown Valley, Soar, with one on 14 June followed by six, including a tandem pair (the second year in a row with evidence of breeding here), on 23 August and two on 1 September; singles at Exeter Canal on 13 June, Stover during 25 June-19 July and Dawlish Warren on 6 September; two males at Clennon Valley Lakes on 8-9 September and at Home Farm, Fremington, on 12 September; and finally a female at East Prawle on 18 September.

2024: Southdown Valley, Soar, on 1 & 16 August and Slapton Ley on 12 & 16 August.

2025: males at Seaton Marshes Borrow Pit on 16 July, Little Haldon on 26 July, Clennon Valley Lakes on 13 August and Soar area on 15 and 25 August.



### **Hairy Dragonfly *Brachytron pratense* (Hairy Hawker)**

*Scarce and local at ditches, ponds and lakes*

Devon holds the most south-westerly outposts of this nationally-increasing species, with strong populations around the Exe marshes, Bovey Basin and the Slapton area. Hairy Dragonfly sometimes wanders away from breeding sites and, in line with the national picture, has colonised new sites in recent years, with the Grand Western Canal and Braunton Burrows being extreme examples.

Around the Exe Estuary, the strongest population extends over the whole of Exminster Marshes, where 60 were counted in 2007 during a systematic survey of the ditches. Other sites include Bowling Green Marsh-Goosemoor (up to four, 1992-2024), Dart's Farm/Lower Clyst Valley (up to 13 since 2005, exuvia at Dart's Farm in 2022), Dawlish Warren (up to three since 1987, with egg-laying in 2006 and 2008), Exeter Canal (including the abutting Alphin Brook, Harpers fishing ponds and Old Sludge Beds DWT Reserve: regular, with occasional breeding evidence and a count of 11 in 2001), Matford Business Park (2003), Matford Marshes (2004) and Matford RSPB (breeding recorded in 2005 and 2009). Rather surprisingly, an egg-laying female was seen at a pond outside the Met Office building in Exeter in 2013.

In the Bovey Basin, where the earliest record comes from 1924, recorded from Decoy CP (up to three since 1993, and at Magazine Pond in 2006 and 2024), Little Bradley Ponds (small numbers since 1992, with egg-laying in 1994 and 2022 and an exuvia in 2024) and Stover CP (the main site, with records since 1982 and regular breeding evidence: daily maximum of 12 exuviae in 2006 and peaks of 10 adults in 2003 and 2025, 13 in 2012, 12 in 2018 and 11 in 2024). One or two at Abbrook Pond (1982), Bovey Heath (2014), Chudleigh Knighton Heath (1995, 1996 and 2009), East Golds (2021), Gallows Cross SUDS ponds (Kingsteignton, 2021-4), Netherexe (and former Key Transport) Pond (1978-96, with egg-laying in 1995), New Bridge settling ponds (2012), the former New Cross Pond (egg-laying in 1992), Rackerhayes (1943-2021, with mating in 1991, larvae in three ponds in 2006 and an exuvia in 2021), West Golds (1992, 1994 and 2005) and Ventiford Ponds (1991, 2014 and 2021 onwards, with 10 males on 22 May 2024 and egg-laying in 2022-24). Several were at Canonteign Falls ponds on 29 May 2025.

In the EDPH, recorded from Aylesbeare Common (small numbers 2001-7 and 2022, including mating in 2001), Bystock Pools (small numbers and breeding evidence regular) and Squabmoor Reservoir (up to three, 2005-19). Elsewhere in east Devon, known in recent years from the Grand Western Canal (since 2003, with up to eight seen and breeding evidence from 2012) and the Otter Marshes (seven north of White Bridge in 2006, five south of White Bridge in 2007), while a territorial male wandered to a West Hill garden pond on 29 May 2009.

In the South Hams, there are outposts at Slapton Ley (up to six since 1978, with breeding evidence in 2005, 2006 and 2010) and Beesands Ley (including egg-laying at an adjacent pond in 2008), and probably at Hallsands.

Following a record from Braunton Burrows on 6 May 2014, a female was seen egg-laying there at SS44843563 on 18 May 2019, the first records for North Devon. There were further sightings there in spring 2022, including a male on 17 May, with one at Partridge Slack on 26 May 2023 and exuviae found there on 2 May 2025. Two at Braunton Marsh on 11 May 2024, at least three each in 2024-25 at Home Farm Marsh and Yelland and one at Godborough on 13 May 2025 were the first to be recorded there.

Flies from late April (earliest 23 April 2009 at Exminster Marshes) to mid-July (latest 5 August 1981, also at Exminster Marshes).

### **Common Clubtail *Gomphus vulgatissimus***

*Vagrant: one record from the River Otter*

One was seen along the River Otter near Tipton St John on 18 July 2017 by a visiting Polish entomologist familiar with this species. Further searches were unsuccessful, although this is an elusive species, after emergence often seen away from water. The species has otherwise not been recorded closer than Gloucestershire, so the origin of this individual is uncertain.

## **Golden-ringed Dragonfly *Cordulegaster boltonii* (Common Goldenring)**

*Fairly common and widespread at or near running water*

Found widely all across the county, from coastal flushes to high moorland streams, and wandering freely away from watercourses. One was on Lundy on 9 September 2009.

An exuvia was found virtually at sea level in Alphin Brook, Countess Wear, in 2009. At Dawlish Warren, also at sea level, in the absence of suitable breeding habitat, there were three records prior to the mid-1990s, but it has since been recorded annually in small numbers.

Flies from late May (earliest 4 May 2007 at Stover) to September/October (latest 18 November at Blanchdown Wood).

## **Orange-spotted Emerald *Oxygastra curtisii***

*Extinct: only record from River Tamar in 1946*

Three specimens (two male) were taken at the River Tamar near Gunnislake by O. G. Watkins in July 1946. There have been no further records here, despite searches, although there is an unconfirmed record of one photographed in the 1960s. The river here, not far above its tidal limit, has calm sections and wooded margins such as those favoured by this species in France, but it is possible that the water quality has been affected by contamination from the old copper mines in the area.

## **Downy Emerald *Cordulia aenea***

*Scarce and very local at wooded ponds and lakes*

Typically only very small numbers are seen at Devon sites, usually territorial males, and egg-laying is observed only rarely. Adults spend much time away from water, around woodland, so seeing them at their breeding sites always makes for a special day. Evidence of breeding comes most often from finding exuviae near the water's edge. Although often in clumps of rushes or Greater Tussock Sedge, emergence can be on trees, even at distance of up to 2m from water and 2m up tree trunks. At Bystock Pools DWT Reserve, exuviae have been found with relative ease on the woodwork of a walkway around the reservoir fringe. After a concerted emergence around mid-May (earliest 4 May 2014 at Bystock), territorial males are seen mainly in late May and June, with smaller numbers into July (latest at Bystock Pools and Squabmoor Reservoir on 29 July 1978).

In the Bovey Basin, which is the main area occupied in Devon, Downy Emerald has been known since a record from 'Bovey Tracey' in 1922. The principle breeding sites have been the former New Cross Pond (until filled in around 1999; maximum of 11 territorial males in 1992), Little Bradley Ponds (originally the older, eastern pond, but also the 'new' pond created in 1992) and Rackerhayes (known from 1978, with up to five territorial males up to 2006). At Little Bradley Ponds, 53 exuviae were found by intensive searching by up to four observers during 12-14 May 2006, giving some idea of the site's productivity (although typically only up to six territorial males patrol the margins). Smaller numbers also breed at Stover CP (presumably in the Lake, although most evidence has come from up to six exuviae

in a day found around the adjacent 'Sue's Pond'; daily maximum no more than four adults) and at Decoy Country Park (where records are rather erratic and shoreline access is very limited, although four, including a mating pair were seen in 2010), including Magazine Pond in 2024.

Elsewhere in the Bovey Basin, recorded from the former Key Transport Pond (now Netherexe) in 1978, and in 1992 there were three territorial males at Abbrook Pond (first here in 1982) and singles at Bradley, Eddison and Rub-a-dub Ponds. Access restrictions at the angling ponds at Abbrook, Eddison, Netherexe and Rackerhayes have hindered surveys in recent years. A male was seen at Ventiford Ponds in 2018, up to three in 2020-23 (including a female in 2021), eight males in 2024 and seven males in 2025. Males were at Bovey Heath on 4 June 2021 and 2 June 2025, presumably wanderers from nearby Stover.

In the EDPH, the species breeds at Bystock Pools DWT Reserve (up to at least six males at the reservoir since 1978; exuviae and egg-laying seen regularly, with at least a dozen exuviae on 21-22 May 2010) and Squabmoor Reservoir (first noted here in 1978; an exuvia found in 2006 and seen egg-laying in 2010; maximum of at least six in 2007 and 2010).

### **Scarlet Darter *Crocothemis erythraea* (Broad Scarlet)**

*Vagrant: one record*

A male at Smallhanger on 16 July 2000 was the fourth record for Britain. A male of the almost identical Oriental Scarlet (*Crocothemis servilia*) emerged from a fish tank in Exeter on 25 June 2010.

### **Broad-bodied Chaser *Libellula depressa***

*Common and widespread at lowland standing waters*

Territorial males are liable to appear at any new or recently-restored pond and even rainwater puddles with bare ground and little vegetation. Often disperses well away from water, where with females and immatures may be found in woodland rides, glades and scrubby areas. Conceted counts at Stover CP have peaked at over 20, mostly territorial males. There are several other records of 20 or more elsewhere, with a maximum of 30 at Chudleigh Knighton Heath in 2006. Flies from mid-April (earliest 11 April 2014 at Westward Ho!) to mid-September (latest 28 October 2018 at Harleston).

### **Scarce Chaser *Libellula fulva* (Blue Chaser)**

*Recent colonist of rivers, drains and pools; may wander*

The full extent of breeding in Devon is not fully known, but likely to be more extensive than at well-known sites in the lower valleys of the Exe, Clyst and Axe and the Grand Western Canal. Systematic surveys and careful checks of rivers, streams, ditch systems pools with lush emergent vegetation at the margins are likely to reveal additional breeding sites, including some of the sites mentioned below.

The first record in Devon was a male by Alphin Brook at Countess Wear, found during a BDS Devon Group meeting on 21 June 2003. However, the first signs of colonisation came in

2005, when 12 were seen along the Grand Western Canal in July, with egg-laying noted there on the 14th. Up to 26 were counted and mating was observed in 2006; in 2007, at least 147 emerged there during 18 May (pictured)-1 June, mainly in the 300m east of Westcott Bridge, with up to 12 seen subsequently; in 2008, at least 53 emerged during 20-30 May, with a subsequent peak of 28 (four egg-laying) on 10 June; 47 were counted on 2 June 2009 and 85 in systematic counts during 31 May-21 June 2010, by which year territories extended along most of the canal, although most were in the central section between Halberton and Ebear. In 2018, 39 were counted in five sections of the canal and 40 in 2021.



In 2007, Scarce Chasers were found along the River Axe around Colyford Bridge (six, including one teneral) and at Whitford, and at Lower Bruckland Ponds (presumably from the nearby River Axe; up to three or four, including a mating pair). In 2008, five were seen at Lower Bruckland and at least six emerged along the River Axe, while at Lower Bruckland numbers had risen to 16 in 2019 and 60+ in 2023 (plus 18 along the Axe). Downriver, a male was seen at a Seaton Marshes drain on 3 June 2011 and singles in the Borrow Pit area during 27 June-11 July 2025. A teneral, presumably from the Axe Valley (at least 3km away) was at Trinity Hill LNR, near Axminster, on 24 June 2016.

The first suggestion of breeding around the head of the Exe Estuary came in 2007 from Exminster Marshes, where the eradication of the invasive alien Floating Pennywort *Hydrocotyle ranunculoides* by the Environment Agency is likely to have produced suitably open ditches: a teneral was found near the M5 on 29 May and six territorial males were along the main drain on 10 June. In 2010, an exuvia was found, egg-laying was observed and 18 (17 territorial males) were present on 22-23 June. The peak count had risen to 46 (43 males) by 2018, 66 (58 males) by 2020, and 92 (88 males) in 2023, mostly still along the main drain (although the extent to which breeding occurs in the ditches across the marshes has not yet been determined). Just 2km to the east, one was found at Dart's Farm fishing ponds, Topsham, on 27 June 2010, mating was seen there in 2016, mating and exuviae in 2017, up to eight males in 2018, mating in 2019, an exuvia in 2022 and more than 20 in 2025. In 2018, four territorial males were found close to Exeter Canal at Harpers Fishing Pools (with individuals there and nearby at Salmon Pool in 2019) and two males in the lower Clyst Valley north of the A376, where there were eight males in 2020; further down the Clyst, an immature male was at Bowling Green Marsh on 30 May 2023. Wandering females were seen at Dawlish Warren on 19 July 2008 and Alphin Brook on 17 June 2012; singles were seen at Powderham Marsh in 2014 and 2022, with six in 2023; the Old Sludge Beds DWT Reserve in 2014, 2017 and 2019 (up to seven), with up to four along Exeter Canal in the same area; however, these last totals were eclipsed by a minimum of 52 (50 males) between Topsham Ferry and Countess Wear in 2023. In 2019, at least two were seen at

Matford Pools in July. Several have wandered into the suburbs of Exminster and Exeter in recent years.

Other singles have been found away from known breeding sites as follows: along the River Culm near Rewe in 2007 (male) and near Hele in 2024 (fresh-looking female), West Charleton Marsh, Kingsbridge, on the late date of 23 July 2010; a fresh-looking female at Buckfastleigh on 23 June 2017; the lower Otter Valley on 30 June 2018; at Clennon Valley Lakes on 25 June 2018, 5-8 July 2019 (male, with a female on 6th), 3 & 22 June 2020 (male), 22 July 2021 (male), 14 June 2022 (at least two males and a female), up to three in 2023; one on 20 June 2024; and up to 10 (including mating pairs) on 17 June-4 July 2025 – the species is presumably breeding here; at Westerland Valley Country Park, Torbay, on 5 June 2019; and at Stover CP (female) on 5 July 2019. At least two were reported at Little Bradley Ponds on 24 June 2020. In 2021, there were records in North Devon from near Winkleigh (15 June), Meeth Quarry DWT Reserve (two males on 26 June and an overmature female on 20 July) and an old female at Lower Tamar Lake on 24 July, and in South Devon from near Woodbury (15 June), the Otter-Tale Valleys at Tawstock (5 June), Otterton (16 June), near Cotley (26 June) and at South Efford Marsh (24 July).

Further signs of range expansion came in 2022, most notably the discovery of 20, including mating pairs, at Braunton Marsh in the spring. Also in 2022, single territorial males were at Little Bradley Ponds and Stover Lake on 2 June, with three males at the latter on 6 June and two males at Squabmoor Reservoir on 3 July (with about six on 18 June 2023 and a single male on 20 May 2025); at Slapton Ley in 2022, 2023 and 2024.

In late May 2023, there were multiple records around the Exe-Clyst marshes following the first at Grand Western Canal on 15<sup>th</sup>, plus two immatures at Meeth Quarry DWT on 29<sup>th</sup> and at least two at Little Bradley Ponds. Dispersing individuals were seen in early June in East Devon at Bystock, Dawlish Warren, River Tale at Escot Park and at Stoneyford, with males at Decoy Lake (3). At Stover, three males were scattered around the lake on 4 June 2023 and eight, including a one egg-laying, on 23 June; none was seen in 2024, but at least 10 (nine males) in May-June 2025. In North Devon, records came again from Braunton Marsh, with six, including a mating pair, on 4 July 2024; a mating pair at Jennetts Reservoir (Bideford); and a male at Canworthy Water, near the Cornwall border, on 22 June. There were four at Dawlish Warren on 22 July 2024, 2-5 at Lower Tamar Lake (including a male with 'mating scars') on 17 July 2024 and at Wooda Lakes and 4 at Jennetts Reservoir in May-July 2024-25.

In 2025, a female was in a Newton Abbot garden on 18 May and at nearby Decoy CP there were at least 10 males on 22 May (after none in 2024), suggesting breeding; 2-5 at Meeth on 23 May 2025; a male was at Escot (R Tale) and at least two males (one with 'mating scars') were at Canonteign Falls ponds on 29 May; one was reported from Little Bradley Ponds again.

Emergence is typically in the second half of May, often resulting in tenerals appearing away from the breeding sites. Adults appear on territory in June and smaller numbers remain until mid-July. The earliest emergence is 6 May 2022 at Exeter Canal and the latest were on 24

July at the Grand Western Canal (2008) and South Efford Marsh (2021), apart from an exceptional record of an immature male at Lower Bruckland Ponds on 1 September 2015.

### **Four-spotted Chaser *Libellula quadrimaculata***

#### *Common and widespread at standing waters*

Widespread, although counts are usually in single figures at most sites. Small numbers are reported from moorland ponds, although 43 exuviae were collected in June 2024 and 35 adults were counted in July 2025 at recently created ponds in the Ockerton Court-Cranmere area of Dartmoor. The highest counts have come from the Bovey Basin, at Little Bradley Ponds (52 on 17 June 2006) and Stover CP (64 on 25 May 2007 and 59 on 4 June 2023). However, 'thousands' occurred in the first four days of June 1963 on Lundy, where many adults emerged from Pondsbury on 15 May 1964; surprisingly, there were no further records from Lundy until 25 May 2023.

At Dawlish Warren, became a regular breeding species after pond excavation in the 1980s but was lost some time in the 1990s, since when there have been one in 2008 and two in 2009, but none since.

Flies from May (earliest emergence 9 April 2011 at Bystock) to August (latest 29 September 2002, also at Bystock).

### **Black-tailed Skimmer *Orthetrum cancellatum***

#### *Common and fairly widespread at lowland standing waters*

This species can be abundant at waters with unvegetated margins and similar basking areas, including around claypits and paths alongside canals.

Maximum counts: 50 at South Huish Marsh on 9 July 2005, Slapton Ley on 11 June 2008 and Stover CP on 22 June 2010 (where also 46 on 24 June 2020); 54 at Southdown Farm, Bolberry, on 26 July 2021; 65 along the Grand Western Canal on 18 June 2010; and over 100 at Lower Bruckland Ponds on 8 July 2006 and at Ventiford Ponds on 3 July 2014.

Flies from mid-May (earliest 30 April 1995 at Dawlish Warren) to August (latest 17 September 2003 near Hallsands).

### **Keeled Skimmer *Orthetrum coerulescens***

#### *Fairly common in wet lowland heath and lower upland bogs and mires*

Keeled Skimmer breeds in boggy mires and seepages and the edges of runnels, streams and ponds in lowland heathland and the lower parts of moorland. It occurs at altitudes of up to 400m on Dartmoor and Exmoor, although is much scarcer at the latter; records from Cramber Pool (430m) and Ockerton Court (2 on 9 July 2025 at 550m) are the highest. Sites typically have abundant bog-mosses *Sphagnum* species.

Devon has many excellent sites for Keeled Skimmer, especially valley mires in Dartmoor, Exmoor and the East Devon Pebblebed Heaths, with sparser records from springline mires in the Blackdown Hills. Heathier Culm grassland sites often hold this species. Between late

June and early August, counts of over 100 have come from Aylesbeare Common (1995 and 2005), Bicton Common (1995 and 2017-20), Doetor Mire (2005), Blackabrook, Lee Moor (2005), Wigford Down (2010) and Vixen Tor Mire (2005), with maxima of 200 at Walla Brook (Riddon Ridge) on 11 July 1999, 200+ at Colaton Raleigh Common in July 2006 and 250 at Smallhanger on 6 August 2006.

Small numbers occur in the Bovey Basin, especially at seepages and drains in clay pits. Water in the ponds here is typically neutral, but the species probably breeds successfully where sufficient *Sphagnum* can establish; it has bred at Chudleigh Knighton Heath (colonised ponds dug in 1990, following the introduction of *Sphagnum*, with a maximum of ten in 1995, but fewer since), Little Bradley Ponds (maximum of 11 in 1995, but none since 2001) and Stover (maximum of five in 2020), and possibly at Ventiford Ponds (maximum of seven males in 2019). There was formerly a good population at Meeth Ball Clay Works, until extraction and pumping ceased and the pits flooded.

Unusually, up to three (including a female) were found in an acidic runnel on the cliff slope at West Soar in 2017-20, with up to 22 in 2021; up to 8 (6 males) were present in 2025.

Immatures are prone to wandering and adults occasionally appear well away from typical habitat, including coastal sites. Recorded at Saltram, Plymouth, in 2010 and 2011, Ford Park Cemetery, Plymouth, in 2023, at Dawlish Warren in 1982 and 1983, while a male was in the dunes there on 24 August 2004 and a worn female was in the same place on 16 August 2010. A female was at Clennon Lakes on 10 July 2024.

Flies from late May (earliest 6 May 2007 at Aylesbeare) to mid-September (latest 15 October 2011 at Vennmills, North Devon).

### **Wandering Glider *Pantala flavescens***

*Vagrant: one record*

One was seen in flight near Otterton on 13 May 2015 by two observers with experience of the species. There have been only three previous British records.

### **Black Darter *Sympetrum danae***

*Fairly common at upland bogs and pools; mainly scarce wanderer in lowlands*

Black Darter is known mainly from acidic pools and mires across Dartmoor and less extensively on Exmoor. Unlike lowland heathland sites in Dorset, Hampshire and Surrey, it breeds rarely and seemingly not very successfully at similar sites in the Bovey Basin, but not the EDPH. National trends suggest that this is one of a small suite of 'northern' species showing a decline during 1970-2019. Along with Emerald Damselfly and Common Hawker, Black Darter has declined in site occupancy (a surrogate for population) in Britain (and on the Continent), which suggests that the decline is related to climate change. However, recent work to restore Dartmoor and Exmoor mires, including the blocking of drains and the creation on many pools, has improved the breeding conditions for this species. Most of the many records from unsuitable breeding sites away from moorland are assumed to have been dispersing from Dartmoor or Exmoor, rather than Continental immigrants.

On Dartmoor, Black Darter occurs widely in blanket and valley mires at 235m ASL or higher, with counts of over 20 in the Burrator catchment and at Gidleigh Common, Gutter Tor Mire (Sheepstor), Quickbeam Hill, Statt's Bridge and Yellowmead (Willsworthy). At extensive peatland restoration around Ockerton Court, up to 39 were seen in 2024 and 97 in August 2025; however, the population across all of the Dartmoor restoration area is likely to run into thousands. Around the fringes of the moor, the species was also known formerly from the Lee Moor clay pits (at 100-150m ASL; maximum 100+ in the Cadover Bridge area, but not since 1999 and not necessarily breeding there) and especially Smallhanger (<200m), where a maximum of more than 100 was counted on 6 August 2006, until the recent extensive loss of habitat due to clay working. A pair, the female egg-laying, was at Smallerhanger, Cadover Bridge, at 210 m ASL in 2024.

Exmoor records have come from Brendon Common (1980-81), Challacombe Common (1999), Furzehill Common (1941) and close to the Somerset border at Badgworthy (2012), Squallacombe (2011) and Lank Combe (2025); it also breeds just over the border near Pinkworthy Pond.

In the lowlands, noted in East Devon from the EDPH at Bystock Pools DWT Reserve in 1977, Bicton Common in 1977, on 2 August 1981 and 2 August 2019, Squabmoor Reservoir on 3 September 1978 and at Aylesbeare and Harpford Commons on 8 September 1999 and 19 July 2010. Further east, recorded at Offwell Woods on 4 August 1993.

In the Bovey Basin, most records have come from at Stover, with two on 10 October 1992, one egg-laying on 21 August 1995, a teneral that flew up from beside a *Sphagnum* area of the 'Puddle Pond' during filming for the BBC's 'One Show' on 29 July 2008, one on 26 August 2017 and a mature male on 18 August 2024. Also recorded at Decoy CP in 1989; singles at Bovey Heathfield on 5 October 2005 and 3 October 2017; single males at Chudleigh Knighton Heath on 29 August 2008 and Little Bradley Ponds on 13 September 2020 (the first there since a record in 1978); Netherexe on 12 September 1978 and Rackerhayes on 1 July 1979. At nearby Haldon Forest, noted at Kiddens Pond on 7 August 1981.

Coastal or near-coastal dispersing records have come from Braunton Burrows (1979), Dawlish Warren (a male on 14 and 23 September 2004), Kenwith Valley NR, Bideford (1 September 1991), Lundy on 29 September 1998 and 21 October 2012 and Torquay (1 September 1930).

Flies from July (earliest 18 June 1995 at Lydford) to early October (latest 12 October 1986 at Holming Beam, Dartmoor, and 21 October 2012 on Lundy Island – the only record for the island).

### **Yellow-winged Darter *Sympetrum flaveolum***

*Vagrant: 12 records*

This erratic migrant to Britain was first recorded in Devon at Woodbury Common in 1928, while there were single records in 1955 at Bovey Heathfield and in 1994 at Marsland NR. During an unprecedented national influx in 1995, singles were at Colaton Raleigh Common on 8 August, Budleigh Salterton on 19 August, a mating pair were seen at Croyde, and at

Berry Head there were five (four males, one female) on 12 August, with two (one male) next day; there were also undated records from Teignmouth and Dawlish Warren (two) and vague records from hectads SX73 (contains Bolt Head and Prawle Point), SX83 (contains Start Point), SX84 (contains Slapton Ley) and SY09 (contains Aylesbeare Common and the upper Otter valley).

### **Red-veined Darter *Sympetrum fonscolombii***

*Very scarce migrant; has bred*

This erratic Continental immigrant has become increasingly frequent in Britain in recent years, with scattered breeding records as the species colonises. Red-veined Darter usually has two generations each year and our climate may not be warm enough to allow this regularly, although the tendency is for bimodal patterns of occurrence, with influxes in June sometimes resulting in breeding and with later emergence in September-October. There have been sporadic breeding records in Devon from five sites (four of them coastal or near-coastal) in at least six years since 1995. These have tended to coincide with national influxes during exceptionally hot weather.

A male that was judged to have emerged from one of the ponds at Dawlish Warren on 21 October 1995 comprised the first evidence of breeding in Devon. Breeding activity was next recorded at South Huish Marsh, with at least six adults and egg-laying seen on 3 August 2003 and a teneral on 9 July 2005, but only erratically since. At Smallhanger, small numbers, most or all males, were recorded in seven years between 1998 and 2009, with a maximum of nine on 11 August 2003, but without any evidence of breeding. However, the species bred around this time some 25 km to the west at fishing lakes near Trerulefoot, Cornwall, and it is not inconceivable that at least some of the Smallhanger individuals had dispersed from here.

Breeding also occurred at Lower Bruckland Ponds in 2006-7: in 2006 egg-laying was observed on several dates between late June and 26 July and up to seven adults, mostly males, were seen; in 2007, a recently-emerged adult and up to six mature adults (one egg-laying) were seen in early June, followed by a male on 24 July and later an exuvia was found on 16 September, an adult on 1 October and a fresh female on 6 October. In 2014, a pair was seen egg-laying at Meeth DWT Reserve on 12 June and single males were seen there in June 2015. In 2019, up to ten were seen in July at both Clennon Valley Lakes and Meeth DWT Reserve, with egg-laying recorded at both sites; both sites had further records up to 2023.

Possibly bred in 2020 at Blackhill Quarry (EDPH), where up to 10 immatures were present in a dry area in June, with 3-4 mature males on territory at a nearby shallow pond on 25 June; however, no exuviae were found then or on 14 September, when a second generation might have been expected. In 2022, present at Southdown Valley, Soar, during 17 June-27 August, with a tandem pair ovipositing on 21 June and 11, including three tandem pairs, on 10 July; four males seen later on 22 September might indicate successful breeding; another good series of sightings in 2023, with evidence of breeding: five males on 24-25 May (clashing with Vagrant Emperors!), two males on 14 and 21 June, eight, including two pairs egg-laying,

on 24 June, and one or two males between 7 July and 1 September, with another pair egg-laying on 23 August. After two singles at Southdown in 2024, present between 15 May and 15 August 2025, including no less than 19 (two pairs egg-laying) on 20 May, with further egg-laying females on 25 June and 11 July.

Other records, generally assumed to be immigrants, are as follows:

1883: recorded at Exmouth (undated).

1903: recorded at Woodbury Common (undated).

1911: collected at Strete on 20 July.

1928: recorded at Grand Western Canal (undated).

1941: collected at Furzehill Pond, Lynton, on 21 and 26 June and 7 July.

1942: recorded at Fernworthy Reservoir on 19 July.

1984: at least six at Beesands and one at Slapton Ley on 17-18 August, one at Dawlish Warren on 22 August and one at Rackerhayes on 27 August.

1992: a male at New Cross Pond 24 May.

1998: singles near Plympton on 20 June and at Dawlish Warren on 12 October.

2000: three at Berry Head on 27 and 30 July.

2001: one at Seaton Marshes on 15 August.

2002: singles at East Prawle on 14 and 19 June, Aylesbeare on 24 June and Start Point on 1 October.

2003: two at Prawle Point on 16 October.

2005: one at Beesands on 16 July and two at Exminster Marshes on 3 and 8 August.

2006: a male at Stover Lake on 10 June; at least six at Mansands on 17-18 June; one at Squabmoor Reservoir on 18 June; a male at Whitepit Quarry (Bovey Basin) on 15 July; three males by the sea at Beesands on 22 July; and single males at Dawlish Warren on 13 August and Otter Marshes on 21 August.

2008: an immature male at Lower Bruckland Ponds on 7 June and a female or immature male on 15 June; a few were seen at South Huish Marsh.

2009: four males at Lower Bruckland Ponds on 7 June and one on 21 June; up to four at South Huish Marsh on 22 July; also two at Berry Head on 4 June, two males at Dawlish Warren on 7 June and one in Plymouth 13 June; a female later at Northam Burrows on 21 September 2009.

2011: a late male at West Soar on 15 November.

2014: a pair egg-laying at Meeth DWT Reserve on 12 June; a male was at Harpers Ponds, Double Locks, on 15 June.

2015: a mature male was on Lundy Island (first record there) on 6 June; males were at Soar on 8 and 15 June; single males were seen at Meeth DWT Reserve on 14 and 29 June; a female at East Soar on 19 June; a male at Dawlish Warren on 21 June; a female at Andrew's Wood on 30 June; and a female was unusually high up (330m) on Dartmoor at Buckland Common on 25 September.

2017: probably at least five different individuals were recorded during 25 May-2 July in the Soar area, plus a male at Dawlish Warren on 1 and 24 June and one at Bovey Heathfield on 5 July.

2018: one at Prawle Point on 13 October.

2019: recorded in early July at Berry Head (three), Bystock Pools DWT Reserve, Dawlish Warren, Stover, Squabmoor and Warren Point, Thurlestone; at least 10, including two pairs egg-laying, at Meeth DWT Reserve on 14 July; at least five on the seaward side of Beesands Ley on 27 July, with possibly more over the Ley; later, a female was at Aylesbeare Common on 18 September.

2020: up to ten immatures away from water at Blackhill Quarry 8-21 June, with three or four territorial males (one pictured) at a nearby shallow pool on 25 June; an immature male at Bovey Heathfield on 6-18 September.

2021: two at Clennon Valley Lakes on 14 June and one in Central Park, Plymouth, on 22 June; up to three males at Soar between 28 June and 8 July.

2022: one flew in off the sea at Dawlish Warren on 27 May and males were on the coastal path east of Budleigh Salterton on 6 July; up to 6 were at Southdown Valley during 17 June-22 September.

2023: males at Exminster Marshes on 16 May and Meeth on 25 May; up to three at Clennon Valley Lakes on 14-15 June, and singles at Dawlish Warren on 29 July and Big Pond, Blackaton Cross, on 14 September.

2024: singles at Soar on 1 June (Warren Barn) and 1 August (Southdown Valley).

2025: males at Braunton Burrows on 2 May and Bicton Common on 20 May; 7 on 16 May at Exminster Marshes; at least ten at Beesands Ley on 15 June; a female near Titchberry, Hartland, on 16 June; and males at Clennon Valley Lakes on 4 July, Berry Head 5 July and Rackerhayes on 9 July.



### **Ruddy Darter *Sympetrum sanguineum***

*Very scarce migrant; has bred*

It is an enigma that this species (and also Brown Hawker) occurs just over the borders of Dorset and Somerset in reasonable quantity (it is very common in the Somerset Levels, for example), but has always been very scarce in Devon.

Larvae were reported from Exminster Marshes on 22 May 1996 (see also under 2002 below) and Sir Arthurs Pill, Braunton Marsh, on 12 September 1996 and Braunton Swanpool on 25 September 1996, respectively. (However, it should be noted that the larvae of this species are notoriously difficult to separate from those of Common Darter.)

Ruddy Darter formerly bred at what appears to have been the main (perhaps only) breeding site in the Bovey Basin: a damp hollow between Rackerhayes pools and Gallows Cross,

Kingsteignton, where 24 adults were present on 3 August 1991, but only two on 17 July 1992; the site was lost when quarried for clay soon after; earlier records from Rackerhayes in 1979 and 1985-86 may refer to this site. At Offwell, East Devon, 'good numbers' were reported before the ponds were virtually cleared of emergent vegetation in the early 1990s: there were at least 20 territorial males in 1994, but only five, including a recently-emerged teneral and one egg-laying, were seen on 14 August 1995, one in 2005, two males in 2008 and a pair egg-laying in 2009; there have been no reports since. An overgrown ditch near the sewage treatment plant at Seaton Marshes held Ruddy Darters until it was dredged in 2013: the records span 1995-2009, with a maximum of 12, including a pair mating and egg-laying, on 22 July 2006.

Seen egg-laying in the Dune Pond at Dawlish Warren in 1994 and four reported there in 1995, but no evidence of breeding since Moorhens took up residence.

On 17-18 July 2006, a total of 44 found at five ponds at Braunton Burrows, with 29 at D Lane Pond (17 males, three teneral, plus four mating pairs), three at I Lane Pond (one male plus a pair mating and egg-laying), six at Snipe Pond (four males plus a pair mating and egg-laying), five males at Vanners Pond and a female at Sheep 1 Pond. However, a male on 11 August 2010 is the only subsequent record.

Other records are as follows:

- 1907: recorded at Dawlish.
- 1924: recorded in hectad SX87 (Newton Abbot area).
- 1949: recorded in hectad SX98.
- 1961: recorded in hectad SX87 (Newton Abbot area).
- 1978: recorded at the old Key Transport Pond on 12 and 18 September and Little Bradley Ponds on 9 July, 5 August and 18 September.
- 1979: recorded at the old Key Transport Pond on 11 July and from Braunton Burrows and New Cross Pond (both undated).
- 1981-2: recorded at Bradley Pond and Little Bradley Pond (undated).
- 1982: recorded at Dawlish Warren.
- 1988: reported from Stover (undated).
- 1989: reported from West Golds, Newton Abbot (undated).
- 1989 and 1993: singles at the Clyst Marshes (undated).
- 1994: one at Exminster Marshes on 15 September.
- 2000: reported from Aylesbeare Common on 19 August.
- 2002: seen at Exminster Marshes from 22 July to 4 October, with a maximum of seven on 29 July – this suggests that Ruddy Darter may breed (or at least may have bred) here – and one or two were at Stover CP during 29 July-28 September.
- 2003: reported from Aylesbeare Common on 3 August and a male at Slapton Ley on 25 August.
- 2004: reported at Chudleigh Knighton Heath on 5 July.
- 2005: different males at Little Bradley Ponds on 18 and 19 July; one at Stover CP on 3 August; three at Riddiford House (near Winkleigh) on 8 August; at least two at Bovey

Heathfield and one near Otterton in August; and one at Dawlish Warren 17-23 October.

2006: a male(s) at Chudleigh Knighton Heath on 18 July, 4 August and 10 September, with two reported on 20 July; one at Sterridge Valley (near Berrynarbor) on 21 July; one at Bystock Pools DWT Reserve on 23 and 25 July; one at Lower Bruckland Ponds on 25 July; one at Dawlish Warren during 26 July-13 August; and one at a farm pond at Deptford Farm, Meddon, on 26 July.

2007: one or two at Beer Quarry, Exminster Marshes, Lower Bruckland Ponds and Seaton Marshes.

2008: an early migrant(?) was at Dawlish Warren in June, a few were seen at several sites in North Devon (including one or two near Beaford) between late July and early September and one was reported in a garden in East Budleigh on 1 September.

2009: one at Smallhanger on 2 August.

2011: reported at Mount Edgcumbe and Slapton Ley on 4 October.

2012: reported at Slapton Ley on 26 August, with two on 18 September (a record of six on 30 October would be exceptionally late for this species).

2013: single males on the coast at Middle Soar on 12 August, Dawlish Warren on 5 October and at Exminster Marshes on the very late date of 23 October.

2014: the first for Lundy Island was on 21 September, clearly a migrant.

2018: one at Lower Bruckland Ponds on 28 July.

2019: singles at Powderham Castle on 10 July and Clendon Valley Lakes on 25 July and several, including mating pairs, at Meeth DWT Reserve 16 July and 26 August.

2020: seen at Meeth on 7 August, one at Trentishoe on 9 August and at least two at Hill Wood, Luxhore, on 1 September.

2021: an immature male at Stover on 4 August.

2022: one at Meeth on 7 August and a presumed migrant male at Soar on 15 September.

2023: a presumed migrant male at Southdown Valley, Soar, on 14 September.

2024: a mature male at Stover on 18 August.

2025: three at Lower Bruckland Ponds on 2 August with one on 5<sup>th</sup>; one at Godborough on 22-25 August.

The records span 5 July (Chudleigh Knighton Heath in 2004) to 23 October (Dawlish Warren in 2005 and Exminster Marshes in 2013), with most in July-August.

### **Common Darter *Sympetrum striolatum***

*Very common and widespread; scarce on high moorland*

This is one of the commonest and most widespread dragonfly species, both in Devon and in Britain, breeding in a wide variety of mainly standing waters. Adults wander widely and are sometimes seen in numbers at coastal locations, even pairs flying in tandem, suggesting immigration. There are at least six records from Lundy Island. At the other extreme, there are records from Dartmoor at 350m (Haytor and Headland Warren), 380m (Shapley Tor) and almost 400m (Devonport Leat at Nun's Cross), while larvae have been seen at Venford Reservoir (about 285m), emergence at Yellowmead, Willsworthy (about 300m) and egg-

laying at Doetor Mire (about 250m). Recent surveys of restored peatland bog pools in the Okement Hill-Cranmere area (500-550m) have revealed small numbers of Common Darters. There is therefore some overlap and potential for competition with Black Darter; climate warming may allow Common Darter to 'invade' these upland waters to the possible detriment of Black Darter.

Counts of 100 or more have come from Bystock Pools DWT Reserve on 22 October 2016; Dart's Farm on 21 August 2005; Little Bradley Ponds on 4 September 1993 and 14 August 1996; Smallhanger in August 2001 and 2005; Stover CP on 9 September 2009 and 10 September 2023; and Upper Tamar Lake on 15 August 1998; the largest count to date is 200 at Seaton Marshes on 15 August 2001. Up to 50 are reported most years on Lundy, where the species has bred.

Common Darter has the longest flight season of any species, especially in Devon, emerging from late May, the earliest being one that emerged at Stover on 14 May 2023. They appear on territories from July and some linger into November or even December, with the latest one on 11 December at Ernesettle.

### **Vagrant Darter *Sympetrum vulgatum* (Moustached Darter)**

*Vagrant: two records*

A female was filmed at the Dune Pond at Dawlish Warren on 6 September 2007, the first in Devon since one was recorded at Torquay on 15 August 1899.

APPENDIX Devon sites for important dragonfly species		Nationally Important Species				Locally Important Species				
Site	Grid Ref.	Southern Damselfly	Scarce Blue-tailed Damselfly	Small Red Damselfly	Scarce Chaser	White-legged Damselfly	Red-eyed Damselfly	Hairy Dragonfly	Downy Emerald	Ruddy Darter
Aylesbeare Common RSPB	SY053906	X		x				?		
Bicton Common	SY040858			X						
Blackaton Cross, Dartmoor	SX573632		X							
Blackhill Quarry	SY031853			X			?			
Braunton Burrows ponds	SS43				?			X		?
Braunton Marsh	SS43				X			X		?
Bystock DWT Reserve	SY033843			X			?	?	?	X
Cadover Bridge area	SX555648		X	X						
Colaton Raleigh Common	SY050868	X?		X						
Combeshead Rewilding (private)	SX384914			X		?				
Dart's Farm, Topsham	SX973885				X		x	?		
Decoy Country Park	SX865702				?		X	?	?	
Emsworthy DWT, Dartmoor	SX74437712			X						
Exeter Canal	SX9291-9686				X	X	X	X		
Exminster Marshes	SX9587/8				X	X	X	X		
Gallows Cross SUDS Ponds	SX861735						X	?		
Gidleigh Common: Moortown Bottom	SX660890	X		X						
Godborough ponds, Bideford	SS437274			X						
Grand Western Canal	ST0113-0718				X		X	X		
Harper's Ponds, Double Locks	SX93099016				?	?	X			
Little Bradley Ponds	SX829778				?		X	X	X	
Lower Bruckland Ponds	SY270930			?	x	?	?			?
Meeth DWT Reserve	SS5308			X	?					?
Prewley Moor, Sourton	SX542909	X		?						
River Axe, Colyford Bridge	SY259926				?	X				
River Clyst and leat, Clyst St Mary	SX970913				?	X				
River Otter and marshes	SY07834				?	X		?		
Slapton Ley NNR*	SX825440				?			X		
Squabmoor Reservoir	SY039841				?		X	?	X	
Stover Country Park	SX834751				X		X	X	X	?
Tor View Moor (private)	SX690918	X								
Venn Ottery Common DWT	SY063921	X						X		
Ventiford Ponds, Teigngrace (private)	SX842749						X	X	?	
Wigford Down, Dartmoor	SX537649		X	X						

X confirmed breeding. x possible breeding. ? no (recent) evidence of breeding.

\* Norfolk Hawker also at Slapton.

